



SCIENCE BASED TARGETS NETWORK  
GLOBAL COMMONS ALLIANCE



LAND

# SBTN Natural Lands Map – Technical Documentation

Version 1.1

Authors: Elise Mazur<sup>1</sup>, Michelle Sims<sup>1</sup>, Elizabeth Goldman<sup>1</sup>, Martina Schneider<sup>1</sup>, Marco Daldoss Pirri<sup>2</sup>, Craig R. Beatty<sup>3</sup>, Fred Stolle<sup>1</sup>, Martha Stevenson<sup>3</sup>

1: World Resources Institute

2: Systemiq

3: World Wildlife Fund – United States

Date last updated: 18 February,  
2025

STEP

3

MEASURE, SET  
& DISCLOSE

## Version History

Version	Update description	Release Date	Effective Dates
1.1	New and improved data inputs. New CC BY-SA 4.0 data license	18 February, 2025	18 February, 2025
1.0	New and improved data inputs	4 September, 2024	4 September, 2024
0.3	Beta version	24 May, 2023	24 May, 2023

# Version 1.1 Change Log

This section describes the changes made between version 1 and 1.1 of the SBTN Natural Lands Maps. The following has been revised for version 1.1:

- New data on cultivated grasslands have been incorporated and old data on livestock density have been removed.
- The methods used to incorporate and harmonize regional data with the natural lands map classes have been revised.
  - In version 1.0 we used all MapBiomias country and biome collections, and in version 1.1 we used only the country collections, with the exception of the Amazon Biome in Guyana, Suriname, and French Guiana.
  - In the new collections for Bolivia, Colombia, Venezuela, Ecuador, and the Amazon, the 'other non-vegetated areas' class was split into 'other non-vegetated *natural* areas' and 'other non-vegetated *anthropic* areas,' which we reclassified to natural bare and non-natural bare, respectively.
  - In version 1.0 we reclassified both the 'non-vegetated areas' (class 22) and 'other non-vegetated areas' (class 25) as natural bare land unless it intersected with UMD cropland or built-up for all countries and biomes. After visual inspection, we determined that the 'other non-vegetated areas' class (25) should always be considered not natural in Brazil. We reclassified this class as non-natural bare unless it intersected with UMD cropland or built-up. We maintained the approach of classifying the 'other non-vegetated areas' in Indonesia, Peru and Chile as natural bare unless it intersected with UMD cropland or built-up.
- We incorporated mining boundaries from Dethier et al. 2023 and Tang et al. 2023
- Some data have been updated to the latest version available. See table below for a full summary of all changes to the input data.
- The description of the datasets, methods, and relevant appendices has been updated to reflect the incorporation of new data and version updates for existing data.
- The Results section (section 3.1-3.3) has been updated with the results and analysis of version 1.1 of the SBTN Natural Lands Map, including validation results for the natural forest class in addition to the binary map. We worked with IIASA to conduct an independent validation of this class specifically due to its relevance for deforestation monitoring.

<b>Dataset name</b>	<b>Description of change</b>	<b>Data category</b>
Global Pasture Watch data	New data	Global data
MapBiomass Brazil Collection 9.0	Replaced Collection 8.0	Regional data
MapBiomass Bolivia Collection 2.0	Replaced Collection 1.0	Regional data
MapBiomass Colombia Collection 2.0	Replaced Collection 1.0	Regional data
MapBiomass Venezuela Collection 2.0	Replaced Collection 1.0	Regional data
MapBiomass Uruguay Collection 2.0	Replaced Collection 1.0	Regional data
MapBiomass Ecuador Collection 2.0	Replaced Collection 1.0	Regional data
MapBiomass Amazon Collection 6.0	Replaced Collection 5.0	Regional data
MapBiomass Chaco Collection 4.0	Removed	Regional data
MapBiomass Atlantic Forest Collection 3.0	Removed	Regional data
MapBiomass Pampa Collection 3.0	Removed	Regional data
United States Geological Survey (USGS) National Land Cover Database, collection 1	Replaced 2019 data, updated year to 2020	Regional data
Tang et al. 2022 Mining Footprint	New data	Global data
Dethier et al. 2023 Alluvial Mining	New data	Global data
Global closed-canopy coconut palm, Descals et al. 2023	New data	Global data
Gridded Livestock of the World 4.0 2020	Removed	Global data

# Version 1.0 Change Log

This section describes the changes made between version 0.3 (beta) and 1.0 of the SBTN Natural Lands Map. The following has been revised for version 1.0:

- New data have been incorporated and some data have been updated to the latest version where available. See table below for a full summary of all changes to the input data.
- The description of the datasets, methods, and relevant appendices has been updated to reflect the incorporation of new data and version updates for existing data.
- The methods used to incorporate and harmonize regional data with the natural lands map classes have been revised. In version 0.3, we reclassified any class in regional datasets with tall woody vegetation (including forest, savannas, woodlands or other mixed classes) as forest in the natural lands map. In version 1.0, we modified this approach to better align with the global AFi definition of natural forest (adopted from FAO) and distinguish natural forest from natural short vegetation in classes that may contain both. For mixed classes in regional datasets that contained both forest and non-forest according to the AFi/FAO definition of forest (such as savanna, woodland, or other mixed classes), we overlaid the UMD tree height data (greater than or equal to 5 meters) to delineate forest versus short vegetation. We continued to classify forest classes in the regional data as natural forest. This change was applicable for the South Africa National Land Cover map and MapBiomias Brazil, Amazon, Chaco, Pampa, Atlantic Forest, Peru, Venezuela, Ecuador, Argentina, and Paraguay. Because many companies setting SBTN targets will also need to comply with the EU Deforestation Regulation, which applies FAO definitions, SBTN is seeking alignment where appropriate. These revisions facilitate better alignment with the EU Deforestation Regulation (EUDR) and provide greater global consistency for definitions used in the map (See Box 1 for more information on EUDR and natural lands map definitions).
- The Results section (section 3.1-3.3) has been updated with the results and analysis of version 1.0 of the SBTN Natural Lands Map.
- Appendix E has been added to document the data licenses of all input data sources.
- Other minor edits to the text to improve clarity.

All new and updated data incorporated in version 1.0 include the following:

Dataset name	Description of change	Data category
MapBiomias Brazil Collection 8.0	Replaced Collection 7.0	Regional data
MapBiomias Amazon Collection 5.0	Replaced Collection 4.0	Regional data
MapBiomias Chaco Collection 4.0	Replaced Collection 3.0	Regional data
MapBiomias Atlantic Forest Collection 3.0	Replaced Collection 2.0	Regional data
MapBiomias Pampa Collection 3.0	Replaced Collection 2.0	Regional data

MapBiomass Indonesia Collection 2.0	Replaced Collection 1.0, updated year to 2020	Regional data
MapBiomass Peru Collection 2.0	New data	Regional data
MapBiomass Bolivia Collection 1.0	New data	Regional data
MapBiomass Colombia Collection 1.0	New data	Regional data
MapBiomass Venezuela Collection 1.0	New data	Regional data
MapBiomass Uruguay Collection 1.0	New data	Regional data
MapBiomass Ecuador Collection 1.0	New data	Regional data
MapBiomass Paraguay Collection 1.0	New data	Regional data
MapBiomass Chile Collection 1.0	New data	Regional data
MapBiomass Argentina Collection 1.0	New data	Regional data
UMD Land Cover	Updated to v2	Global data
United States Geological Survey (USGS) National Land Cover Database, Dewitz, J. 2021	New data	Regional data
European Space Agency WorldCereal Active Cropland, Van Tricht, K., et al. 2023	New data	Global data
Digital Earth Africa Cropland Extent 2019	New data	Regional data
Global closed-canopy coconut palm, Descals et al. 2023	New data	Global data
Gridded Livestock of the World 4.0 2020	Updated year to 2020	Global data
LUCAS New Zealand Land Use Map 2020	Updated year to 2020	Regional data

# Contents

- Version 1.1 Change Log ..... 2
- Version 1.0 Change Log..... 4
- 1. Introduction ..... 7
- 2. Data and Methods.....10
  - 2.1. Definitions.....10
  - 2.2. Datasets..... 16
    - 2.2.1. Land Cover classes .....17
    - 2.2.2. Supplementary global data ..... 19
      - Forests ..... 20
      - Short Vegetation and Bare ..... 21
      - Wetlands ..... 21
      - Cropland..... 22
      - Built-up..... 22
    - 2.2.3. regional data ..... 22
  - 2.3. Methods..... 26
    - 2.3.1. Methods for creating and combining map classes for global data .....26
      - Pre-processing ..... 29
      - Forests ..... 29
      - Short vegetation and bare ..... 30
      - Water..... 30
      - Built-up..... 30
      - Combining map classes ..... 30
    - 2.3.2. Methods for incorporating regional data.....30
      - Regional data using single or select classes ..... 31
      - Regional data using all classes ..... 31
    - 2.3.3. Methods for creating the final map ..... 32
    - 2.3.4. Validation methods..... 32
- 3. Results ..... 34
  - 3.1. Map Results ..... 34
  - 3.2. Validation and accuracy ..... 37
  - 3.3. Comparison with existing data ..... 38
- 4. Limitations ..... 39
- 5. Future Work ..... 43
- 6. Data Availability ..... 43
  - Acknowledgements ..... 43
- Appendix A: MapBiomass Reclassification..... 44
- Appendix B: South Africa National Land Cover Reclassification..... 49
- Appendix C: LUCAS New Zealand Land Use Reclassification ..... 55
- Appendix D: SDPT v2.0 Data Sources ..... 58
- Appendix E: Data Source Licenses..... 66
- References..... 68

# 1. Introduction

Natural lands are being lost and degraded at unprecedented levels (IPBES, 2019). Three-quarters of land and two-thirds of oceans have been significantly impacted by humans through pollution, urban expansion, conversion to crop or livestock production, intensive logging in natural forests, unsustainable fishing practices and other activities. The rate of species extinctions is also accelerating, with some experts warning that a sixth mass extinction may be under way. Wildlife populations have decreased by 69% since 1970 (WWF, 2022), and projections estimate that three-quarters of today's animal species could go extinct within 300 years (Barnosky et al. 2011).

Climate change is intertwined with natural land loss. According to the Intergovernmental Panel on Climate Change (IPCC 2019), emissions from the agriculture, forestry and other land use sectors contribute 23% of all anthropogenic emissions. In 2018, the IPCC warned that to avoid the worst impacts of climate change, global warming must not exceed 1.5°C above pre-industrial temperatures. To achieve this, greenhouse gas (GHG) emissions must halve by 2030, and drop to net-zero by 2050. A variety of stakeholders within agriculture, forestry, and other land use sectors around the world have met this call to action with ambitious pledges to reduce and eliminate their share of GHG emissions. However, the sector-wide transformations needed to achieve net-zero by 2050 require coordination and guidance on how to do this effectively and efficiently.

To achieve this goal, voluntary initiatives have emerged to help companies better understand their impact on nature and emissions contributions and determine a plan for reducing or eliminating this impact. The Science Based Targets Network (SBTN) is one such voluntary initiative. It builds on the progress of the Science Based Targets initiative (SBTi) which enables companies to set science-based greenhouse gas emissions reductions and net-zero targets. SBTN is a network of international environmental nonprofit organizations, international agencies and mission-driven entities developing methods and resources for science-based targets (SBTs) for nature for companies, and forthcoming methods and resources for science-based targets for both climate and nature for cities. SBTN's goal is for the world's major companies and cities to have adopted science-based targets and taken action for climate, water, land, ocean and biodiversity by 2025. This will form a key part of progress towards meeting the commitments of the Paris Agreement, The Kunming-Montreal Global Biodiversity Framework, Land Degradation Neutrality, and the Sustainable Development Goals. Additionally, the European Union Deforestation Regulation (EUDR), adopted in May 2023, requires companies that sell key deforestation-driving commodities<sup>1</sup> in the EU to conduct due diligence and verify that their products are deforestation or degradation-free as of 2020, meaning they are sourced from land where no deforestation or forest degradation has occurred since 2020. This new regulation is the first legal application of no deforestation or conversion goals.

---

<sup>1</sup> Cattle, cocoa, coffee, palm oil, soy and rubber, as well as derived products such as beef and chocolate are required to be deforestation-free, while wood and derived products such as furniture are required to be degradation-free.

### Box 1: Definitions used in the SBTN No Conversion of Natural Ecosystems Target and the EU Deforestation Regulation

The SBTN No Conversion of Natural Ecosystems Target uses the Accountability Framework initiative (AFi) definitions and operational guidance to define natural ecosystems, deforestation, and conversion. Although the EU Deforestation Regulation (EUDR) and AFi generally take a similar approach in how they address deforestation, there are some key differences.

AFi and EUDR both adopt the FAO definition of forests, with the exception of rubber, which is considered forest under FAO and agriculture under EUDR. AFi further distinguishes ‘natural forests’ from ‘tree plantations’, while EUDR similarly distinguishes natural from planted or plantation forests using FAO definitions of ‘plantation forest’, ‘planted forest’, ‘primary forest’, and ‘naturally regenerating forest’. However, a key difference between the two frameworks is how they define deforestation: while AFi defines deforestation as the conversion of *natural forest* to agriculture, other non-forested land use, tree plantations, or other forms of severe degradation, EUDR defines deforestation as the conversion of *all forest* to *agricultural use*. Meanwhile, EUDR considers conversion of primary forest into plantation forest, planted forest, or other wooded land as ‘forest degradation’, as well as conversion of naturally regenerating forest into plantation forest or other wooded land as ‘forest degradation’.

In practice, this means that conversion of plantation forest to agriculture would be considered deforestation under EUDR, but would not be considered deforestation under AFi. Meanwhile, conversion of natural forest to plantation forest would be considered deforestation under AFi, while it would be considered degradation under EUDR. Both frameworks consider conversion of natural forest to agriculture as deforestation. For more information on how the two frameworks compare, see the AFi Operational Guidance on Applying the Definitions Related to Deforestation, Conversion, and Protection of Ecosystem (AFi 2023).

Because the natural lands map aligns with AFi definitions in order to facilitate target-setting under SBTN, its primary focus is mapping natural lands (including natural forest), as it is only conversion of natural forest that is considered deforestation under AFi. In the natural lands map, tree plantations are included with tree crops and other tree cover found in predominantly urban or agricultural land uses in the ‘Non-natural tree cover’ class. Users should be cautious of these differences when seeking to use the map as a resource to support monitoring under EUDR.

SBTN’s Land Hub has developed a version 1 of its voluntary corporate target setting methodology. These targets address land occupation and land transformation and are intended to be quantifiable targets for companies to set as they begin engaging with SBTN. Besides their specific contributions to achieving global goals for nature, the design of these three land targets is intended to address the potential trade-offs that might result from different corporate response options to the negative impact to nature in their supply chains. All three targets work in conjunction to prevent negative consequences from one of the targets alone. The targets are:

1. No conversion of natural ecosystems
2. Land footprint reduction
3. Landscape engagement



The first among these land targets, no conversion of natural ecosystems, recognizes the value of all terrestrial ecosystems. Until recently, commitments were primarily focused on forests and achieving zero-deforestation targets (Taylor et al. 2022). However, this ignores other valuable and vulnerable natural ecosystems. For example, grasslands and other short vegetation ecosystems like shrublands are one of the largest biomes on Earth and are rich with biodiversity, yet they are particularly susceptible to conversion because they are easier to clear than forests. Wetlands are vulnerable to development, despite their critical role in providing habitat, improving water quality, and preventing floods. All natural ecosystems store and sequester carbon, support biodiversity, regulate the climate, filter air and water, protect communities from flooding, regulate against diseases and pests, and provide food, medicine, fuel and shelter for Indigenous People and local communities.

Preventing the conversion of natural ecosystems starts with knowing where natural lands exist by delineating them on a map. This Map:

- Provides companies and other stakeholders with a baseline from which to estimate their conversion of natural lands from 2020 with their current production unit or sourcing area data.
- Provides a baseline for independent groups to monitor conversion of natural lands.
- Allows companies to set no conversion of natural ecosystems targets under SBTN.
- Is developed based on a specific definition of natural ecosystems and lands. If using this map outside of the SBTN context, users should compare definitions.
- Provides a 2020 baseline that is agreed upon by a broad membership of organizations, including those of the SBTN Land Hub and the Accountability Framework Initiative (AFi).
- Like all maps, contains errors, and additional validation, including with satellite imagery, field visits, or cross-checking with other data sources may be useful.

The SBTN Natural Lands Map does **not**:

- Contain time-series data that may be useful for monitoring conversion.
- Quantify the area of natural and non-natural lands because of known overestimation of natural lands.
- Supplant existing research and biophysical mapping and analysis on ecosystem science.
- Define ecosystems and/or working lands.
- Assess the importance of the natural land for biodiversity.
- Assess the quality of ecosystems.
- Represent an unbiased map of natural lands – the conservative approach used overestimates the extent of natural lands, and while remote sensing data, on which the map is based, can provide powerful insights, additional field work should be used for validation and to understand local dynamics. Caution should be used if calculating areas with the SBTN Natural Lands Map.

This technical note outlines the methods, results and limitations of the version 1 2020 baseline SBTN Natural Lands Map. The map will be used by companies setting an SBTN No Conversion of Natural Ecosystems Target to understand the baseline area of natural lands for monitoring progress toward zero-conversion of natural ecosystems going forward.

## 2. Data and Methods

Our approach for identifying natural lands across the world was to combine the best available global spatial data on land cover and land use into a single harmonized map at 30 meter resolution circa the year 2020. We aligned our definitions and approach to the extent possible with the Accountability Framework initiative (AFi) definitions (AFi 2024) of natural ecosystems and AFi Operational Guidance on Applying the Definitions Related to Deforestation, Conversion, and Protection of Ecosystems (AFi 2023), recognizing the limitations of what can be directly mapped with earth observation data and relying on proxies to operationalize these definitions based on existing land cover/land use and supplementary data. We assessed and selected the land cover and land use data that were best suited for distinguishing between natural and non-natural land, using additional data where necessary and possible.

While a global approach to mapping natural lands can help produce consistent, comparable results, local ecosystems are not always well represented with global data. For example, global forest data calibrated at broad regional scales can underestimate dry forests (Bastin et al. 2017). Where possible, the natural lands map incorporates and prioritizes regional data to better represent local ecosystems.

### 2.1. Definitions

Science Based Targets Network (SBTN) adopted the AFi definitions of natural ecosystems and forests, which were used as guidance for developing the map. AFi defines a natural ecosystem as “one that substantially resembles - in terms of species composition, structure, and ecological function - what would be found in a given area in the absence of major human impacts,” and can include managed ecosystems as well as degraded ecosystems that are expected to regenerate either naturally or through management (AFi 2024). Because species composition and ecological function cannot be directly mapped with earth observation data, our approach operationalizes AFi definitions using proxies based on available data that align with AFi guidance to the extent possible. We used AFi Operational Guidance in Applying Definitions (AFi 2023) to guide our development of proxies (see Table 1).

While natural forests are of course part of natural ecosystems, a detailed forest definition is also provided by AFi, as adopted from the Food and Agriculture Organization of the United Nations (FAO). At a global scale, forests are defined as “land spanning more than 0.5 hectares with trees higher than 5 meters and a canopy cover of more than 10 percent, or trees able to reach these thresholds in situ. It does not include land that is predominantly under agricultural or other land use” (AFi 2024). At regional and local scales, AFi guidance states that “quantitative thresholds (e.g., for tree height or canopy cover) established in legitimate national or sub-national forest definitions may take precedence over the generic thresholds in this definition” (AFi 2024). For this reason, forest classes in the regional data sources were accepted, even when height or canopy cover thresholds were not specified. However, for classes in regional data sources that contain both forest and non-forest vegetation as defined by AFi/FAO (such as savannas, woodlands, or other mixed classes), we used additional data to apply the tree height and minimum mapping unit thresholds used in the AFi definition.

Table 1. Afi operational guidance and description of how it was used to develop the mapping approach. Specific data and methods used are described in section 2.2 and 2.3.

Afi classification	Attributes and descriptions in Afi operational guidance, with <i>italics indicating elements that cannot be directly mapped with remotely sensed data</i>	Description of how Afi guidance was used to determine operational proxy	Limitations of proxy
<b>Natural Forest</b>			
<b>Unmanaged or minimally managed natural forest</b>	<ul style="list-style-type: none"> <li>Unmanaged or minimally managed natural forest, including with some human impacts.</li> </ul>	<p>We used the definition of forests (tree cover greater than 5 meters in height and more than 0.5 hectares) and the process of elimination to map natural forests by labeling plantations, planted forest or tree crops as non-natural.</p>	<ul style="list-style-type: none"> <li>May include tree plantations and tree crops when data is not available for specific regions or crop types.</li> </ul>
<b>Managed natural forest</b>	<ul style="list-style-type: none"> <li>Forests that are managed for harvest or services <i>in a way that maintains most of the key elements of ecosystem composition, structure, and function over time.</i></li> <li>Forests undergoing selective harvest <i>where high value species are planted or promoted.</i></li> </ul>	<p>Where regional data were used, definitions of forest may not specify a height or canopy density threshold.</p> <p>When evaluating supplementary data or regional data, any class name or description that included “natural”, “native”, “naturally regenerating”, or “secondary” were considered natural.</p>	<ul style="list-style-type: none"> <li>May include severely degraded forest.</li> <li>May include areas under shifting cultivation, regardless of the length of fallow period or impact.</li> <li>May include tree cover within agricultural mosaics, regardless of whether it is under agricultural use.</li> </ul>
<b>Regenerated natural forest</b>	<ul style="list-style-type: none"> <li>Forests that have regrown <i>and now have ecosystem composition, structure and function similar to forest native to the site.</i></li> <li>Regrowth of <i>native</i> vegetation for several years after agricultural abandonment.</li> <li><i>Plantings of diverse native tree species through management for ecosystem restoration.</i></li> </ul>		<ul style="list-style-type: none"> <li>May include agroforestry, regardless of intensity or whether crops are grown under forest canopy.</li> </ul>
<b>Non-permanent or low-intensity cultivation</b>	<ul style="list-style-type: none"> <li>Permanent, semi-permanent, or shifting cultivation that causes little</li> </ul>		

<p><b>within a natural forest</b></p>	<p>disturbance of the canopy <i>and retains a high proportion of species and main attributes of the forest's structure and function.</i></p> <ul style="list-style-type: none"> <li>• Swidden cultivation in small, isolated patches harvested for short periods and then left fallow.</li> <li>• Low-intensity forest farming such as some rustic coffee and rubber agroforestry systems <i>under forest canopy.</i></li> </ul>		
<p><b>Natural non-forest ecosystem</b></p>			
<ul style="list-style-type: none"> <li>• Largely “pristine” natural ecosystems that have not been subject to major human impacts in recent history.</li> <li>• Regenerated natural ecosystems that were subject to major impacts in the past (for instance by agriculture, livestock raising, tree plantations, or intensive logging), but where the main causes of impact have ceased or greatly diminished <i>and the ecosystem has attained species composition, structure and ecological function similar to prior or other contemporary natural ecosystems.</i></li> <li>• Managed natural ecosystems (including many ecosystems that could be referred to as “semi-natural”) <i>where much of the ecosystem's composition, structure, and ecological function are present, including native grasslands or rangelands that are, or have historically been, grazed by livestock.</i></li> <li>• Natural ecosystems that have been partially degraded by anthropogenic or natural causes (e.g. harvesting, fire, climate change, invasive species, or others) but where the land has not been converted to another use <i>and where much of the ecosystem's composition, structure, and ecological function remain present or are expected to regenerate naturally or by management for ecological restoration.</i></li> </ul>	<p>We used the process of elimination to map natural short vegetation and bare ground by labeling all short vegetation and bare ground with high densities of ruminant livestock, cropland, or tree crops as non-natural.</p> <p>When evaluating regional or supplementary data, any class name or description that included “natural”, “native”, “low-intensity grazing”, “secondary”, or “naturally regenerating” was considered natural.</p>	<ul style="list-style-type: none"> <li>• May include unstocked forest.</li> <li>• May include areas used for livestock grazing.</li> <li>• May include fields used for various purposes, including recreation or agricultural activities.</li> <li>• May include areas under shifting cultivation, regardless of the length of fallow period or impact.</li> <li>• May include severely degraded non-forest ecosystems.</li> </ul>	

<ul style="list-style-type: none"> <li>• Grasslands, savannahs, wetlands, and other areas that are not recently transformed or intensively managed, <i>and maintain much of the ecosystem's structure, composition, and function.</i></li> <li>• Includes many traditional pastoral systems and <i>well-managed</i> livestock grazing on native vegetation.</li> </ul>			
<b>Tree plantation</b>			
<ul style="list-style-type: none"> <li>• Eucalyptus or rubberwood plantations.</li> <li>• Monocultures of temperate or boreal species where such monocultures would not have naturally existed.</li> <li>• Monoculture and/or even-aged management where such management does not approximate the spatial and temporal dynamics of a natural forest ecosystem.</li> <li>• All or a substantial portion of planted trees are exotics.</li> <li>• <i>Regular herbicide or pesticide usage.</i></li> </ul>	<p>We used available data on tree plantations, wood fiber or timber plantations, and planted forests.</p> <p>When evaluating local and supplementary data, any class name or description that included “plantation” or “planted” were considered non-natural.</p>	<ul style="list-style-type: none"> <li>• Tree plantations are not mapped comprehensively for all regions. Therefore, tree plantations may be mapped as natural forest.</li> <li>• Planted forests are not mapped comprehensively for all regions, and data on forest management is extremely limited and may not contain sufficient detail on management intensities. Therefore, some monoculture/even-aged management may be mapped as natural, and likewise some semi-natural planted areas may be mapped as non-natural, even if meeting Afi criteria for natural forest.</li> </ul>	
<b>Agriculture</b>			
<b>Permanent smallholder agriculture for local consumption and trade; permanent agriculture for commodity production</b>	<ul style="list-style-type: none"> <li>• Cultivation of temporary or annual crops that have a growing cycle of one year or less.</li> <li>• Cultivation of permanent or perennial crops that have a growing cycle of more than</li> </ul>	<p>We used available data on:</p> <ul style="list-style-type: none"> <li>• Cropland</li> <li>• Tree crops</li> <li>• Specific crop types</li> <li>• High ruminant density</li> </ul>	<ul style="list-style-type: none"> <li>• Tree crops are not mapped comprehensively for all crop types or for all regions. Therefore, some tree crops may be mapped as natural forest.</li> <li>• Due to the dynamic nature of shifting</li> </ul>

	<p>one year, including tree crops.</p> <ul style="list-style-type: none"> <li>• Cultivation of permanent or temporary meadows or pastures, for example by planting of non-native grasses and/or by agricultural management practices such as irrigation or fertilization.</li> <li>• Raising of livestock on land characterized by severe and sustained degradation.</li> <li>• Buildings, animal feeding operations, and other farm infrastructure.</li> <li>• Temporarily fallow land.</li> </ul> <p>For boundary cases, may include:</p> <ul style="list-style-type: none"> <li>• Intensification of swidden agriculture in which patches become larger, cultivation periods longer, fallows shorter.</li> <li>• <i>Cultivation leads to significant and long-term change in ecosystem composition, structure, and function.</i></li> </ul>	<p>areas</p> <p>When evaluating regional and supplementary data, any class name or description that indicated “mixed agriculture”, “agricultural mosaic”, “pasture”, “high-intensity grazing”, or “cultivated” was considered agricultural use and thus non-natural.</p>	<p>agriculture, it is often not included in data on cropland. Therefore, shifting agriculture, regardless of length of fallow period or impact, may be mapped as natural.</p>
<b>Severely degraded land</b>			
<ul style="list-style-type: none"> <li>• Land formerly meeting the definition of a natural ecosystem (either forest or non-forest) <i>that has experienced severe and sustained degradation that alters ecosystem composition, structure and function to the extent that regeneration to a prior state is unlikely.</i></li> </ul> <p>Degraded natural ecosystems (including forests) are generally presumed to be natural</p>	<p>Because ecosystem composition and function cannot be directly mapped with remotely sensed data, we only classified severely degraded areas as non-natural if they</p>	<ul style="list-style-type: none"> <li>• Severely degraded forest patches or other ecosystems within agricultural areas may be mapped as natural if they are not specifically mapped as agriculture by</li> </ul>	



<p>ecosystems unless:</p> <ul style="list-style-type: none"> <li>• The land is managed for uses other than natural ecosystem.</li> <li>• <i>Due to severe or sustained degradation, the ecosystem is not able to regenerate much of its prior ecosystem structure, composition, and ecological, biophysical, and cultural functions naturally and/or through assisted regeneration.</i></li> </ul>	<p>were mapped by existing data as within an agricultural or built-up extent.</p>	<p>existing data.</p> <ul style="list-style-type: none"> <li>• In general, severely degraded ecosystems may be mapped as natural.</li> </ul>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------

Note: AFi classification, descriptions, and attributes are from AFi definitions and Tables 1 and 2 from AFi Operational Guidance on Applying the Definitions Related to Deforestation, Conversion, and Protection of Ecosystems (AFi 2023).

We also considered AFi’s conversion definition in anticipation of the relevance of this natural lands map for monitoring purposes, which includes “a change to another land use or profound change to composition, structure, or function” (AFi 2024). Conversion can happen regardless of whether or not the change was legal. In this technical note we do not map or monitor conversion.

Additional natural land cover classes beyond forests were included in the map: short vegetation, which includes grasslands and shrublands, water, snow/ice, bare land, and wetlands (Table 2). In the absence of specific definitions for these ecosystems from AFi, we relied on definitions from available data sources. Here, short vegetation is defined as areas of land with vegetation shorter than 5 meters, and can include areas of land dominated by grass or shrubs. Water is defined as surface water present 20% or more of the year, outside of wetlands. Snow and ice include any permanent snow and ice. Wetlands are transitional ecosystems with saturated soil that can be inundated by water either seasonally or permanently, and can be covered by short vegetation or trees. Bare land is defined as areas with exposed rock, soil, or sand with less than 10% vegetated cover. Table 2 includes examples of the types of ecosystems which may be included under these broad land cover classes.

Table 2. Examples of ecosystem types that may be included under the map’s natural land cover classes.

Natural land cover class	Class definition	Ecosystem examples
Forest	Areas with tree cover greater than or equal to 5 meters in height spanning more than 0.5 hectares.	Rainforests, dry forests, montane rainforests, heath forests, temperate forests, boreal forests, some types of savannas or woodlands.
Short vegetation	Areas of land with vegetation shorter than 5 meters, including areas of land dominated by grass or shrubs.	Grasslands, shrublands, heathlands, steppes, vegetated deserts and semi-deserts, some types of savannas or woodlands.
Wetlands	Transitional ecosystems with saturated soil that can be inundated by water either seasonally or permanently, and can be covered by short vegetation or trees.	Peatlands, mangroves, inland, coastal, saline, freshwater, brackish.

Water	Surface water present 20% or more of the year, where water is the dominant class.	Rivers, lakes, coastal inlets, bays, lagoons.
Snow/Ice	Areas covered by permanent snow or ice.	Glaciers, perennial snowfields.
Bare land	Areas with exposed rock, soil, or sand with less than 10% vegetated cover.	Sparsely-vegetated deserts, lava flows, screes, alpine rocky outcrops, sandy shorelines.

*Note: The ecosystem examples included in this table are not an exhaustive list of all ecosystems included within each land cover class, but are illustrative examples of some types of ecosystems which may be included. Land cover classes are defined based on the biophysical presence and coverage of certain types of vegetation or landforms, and thus a similar type of ecosystem in different regions may fall into different land cover classes depending on the biophysical characteristics present. Please note that in cases where regional data was incorporated, there may be inconsistencies in how some land cover classes are defined (e.g. with regard to tree height threshold for forests, etc.).*

## 2.2. Datasets

The natural lands map combines data collected from a variety of sources that were assessed for quality and met certain criteria (Table 3). Additionally, all data – including regional data sources – were subject to a visual inspection using high resolution imagery as an added assurance that the land cover classes selected aligned to the extent possible with AFI definitions.

Table 3. Selection criteria for natural lands map data.

Licensing	Data included in the map should be publicly accessible and licensing should allow for a wide variety of uses.
Resolution	Data included in the map should ideally have a spatial resolution of 30 x 30 meters or higher. If no 30 x 30 meter data are available, coarser resolution data can be included to fill any data gaps. Vector data are also suitable for inclusion if high resolution raster data is not available.
Timescale	Data included in the map should be as close to the year 2020 as possible, but not after it.
Accuracy	Data included in the map should have robust user and producer accuracy scores when available. When using a specific class within a dataset, we looked at individual class accuracy. Accuracy was considered, along with the other selection criteria, when comparing among available data.
Definitions	Class definitions are aligned with our mapping needs.
Coverage	Data included in the map should have a global extent to ensure all geographies have coverage. However, regional data that meet the other requirements outlined in this table, and which define land cover classes and natural ecosystems in a way that is aligned with our mapping needs, should take precedence over global sources.



First, we assessed and selected global land cover data to establish the base land cover classes in the natural lands map. Because most global land cover maps define vegetated classes based on the biophysical presence of vegetation types and do not contain information on the degree of human impact or other characteristics that can be used to delineate natural ecosystems according to the AFi definition, we evaluated additional supplementary datasets to distinguish natural and non-natural lands for specific land cover classes.

2.2.1. LAND COVER CLASSES

The land cover classes included in the map are largely drawn from two maps of global land cover for 2020: (a) WorldCover, a 10 meter resolution dataset created by the European Space Agency (ESA) (Zanaga et al. 2021), and (b) Global Land Use and Land Cover Change v2, a 30 meter resolution dataset created by the Global Land Analysis and Discovery Lab at the University of Maryland (UMD) (Hansen et al. 2022; Potapov et al. 2022). The classes in the UMD data were mapped independently and then overlaid into a composite; for this work we used the independent classes for forest, water, cropland, built-up (v2), wetland, vegetation fraction, and snow/ice. Both UMD and ESA data share a similar classification scheme, and were compared to decide which land cover classes from each product were most appropriate for our map (Table 4A and 4B).

Table 4A. Breakdown of land cover classes and measures of user accuracy (UA) and producer accuracy (PA) as reported in their technical documentation. Bold indicates that the data were included in the natural lands map.

Map Class	ESA	UA   PA	UMD	UA   PA
Forests	Trees	80.8   89.9	<b>Forest</b>	94.6   94.8
Short vegetation	<b>Shrubland</b>	38.6   44.1	Short Vegetation	N/A
	<b>Grassland</b>	69.3   76.7		
Wetlands	<b>Herbaceous Wetland</b>	27.8   40.6	<b>Wetland Short Vegetation</b>	52.4   59.6
	Mangroves	68.6   51.5	<b>Wetland Forest</b>	
	Open Water	88.5   85.0	<b>Permanent Water</b>	98.8   86.1
Non-Natural	Cropland	81.1   76.7	<b>Cropland</b>	88.5   86.4
	Built-up	67.7   67.9	<b>Built-up</b>	63.7   39.1
Other	<b>Barren/Sparse Vegetation</b>	87.5   81.4	<b>Bare</b>	N/A
	Snow and Ice	93.9   97.0	<b>Snow/Ice</b>	63.7   39.1

Table 4B. Summary of comparison between ESA WorldCover and UMD Land Cover data, notes from visual inspection of the data, and decision-making process.

Map Class	Comparison Description	Decision
Forests	The UMD forest class had higher accuracy than the ESA WorldCover tree class.	UMD
Short Vegetation	The UMD short vegetation class was made by clipping out other classes from a global vegetation fraction dataset, and therefore is not intended to stand on its own. We defaulted to the ESA WorldCover shrubland, grassland, and herbaceous wetland classes which were independent classes in the WorldCover map. Although the UMD data were ultimately selected to delineate wetlands, the ESA WorldCover herbaceous wetland class was included in the short vegetation class because it contains areas with vegetative cover, which are included in our short vegetation definition.	ESA
Wetlands	The UMD wetlands data benefit from a higher accuracy score as well as a general “wetland forest” class.	UMD
Cropland	The UMD cropland accuracies were higher than those of the ESA WorldCover cropland class. While UMD’s cropland class is older (2016 - 2019), it classifies areas which had crops during any of those four years as cropland, allowing for a fallow period. By definition, the ESA cropland class does not include cropland that was in fallow in 2020. Because we aim to include all areas used for crop production – including both temporarily fallow and cultivated cropland – in our cropland class, the UMD data better suited our needs.	UMD
Built-Up	While the UMD accuracy scores are lower than ESA WorldCover, we selected the UMD built-up class because its definition includes any pixels that contain man-made constructions or surfaces, including lower density built-up areas such as airports and suburban neighborhoods. The ESA built-up class includes only pixels covered by buildings, roads, and other man-made surfaces, while excluding parks, sports facilities, and other urban green spaces. The UMD built-up class therefore includes more areas which fall under our definition of non-natural.	UMD
Bare Land	The ESA barren/sparse vegetation class definition includes areas with exposed rock, soil, or sand with less than 10% vegetated cover, whereas the UMD bare class is derived from the global vegetation fraction dataset and includes lands with less than 7% vegetated cover. Therefore, the ESA barren/sparse vegetation class classifies a larger area as bare land, including areas such as alpine rock faces, whereas these areas are classified as short vegetation with the UMD data. We therefore combined the UMD and ESA classes with a union to capture the extent of both to provide broader coverage of this class.	ESA and UMD
Snow and Ice	Both UMD and ESA snow and ice classes had high accuracies, however upon visual inspection, ESA seemed to overestimate snow and ice, leading us to use the UMD class.	UMD
Water	The water class in the UMD data had a higher User’s Accuracy, and was adjustable based on the percentage of the year water was present.	UMD

Overall, we found that the UMD Land Cover data were a better fit for the map for most classes, with the exception of the short vegetation and bare classes. While the ESA data benefit from having a higher spatial resolution and therefore more precise data, accuracy metrics were generally lower. Further, we wanted to choose data where the spatial resolution was as consistent as possible; global data with a 10 meter spatial resolution are uncommon and would make resampling difficult.

2.2.2. SUPPLEMENTARY GLOBAL DATA

To distinguish natural from non-natural lands in the land cover classes that contain both, we incorporated additional global data into the map (Table 5). While both the ESA and UMD Land Cover data include non-natural classes (cropland and built-up area), the other land cover classes selected from both the ESA and UMD Land Cover data include areas that do not adhere to AFI’s definition of a natural ecosystem.

Table 5. Summary of supplementary global data used to delineate natural and non-natural lands by land cover class.

Classification	Land cover class	Dataset name	Resolution	Year	Reference
Non-natural	Tree cover, short vegetation	Spatial Database of Planted Trees (SDPT), version 2.0	Varies	Varies	Richter et al. 2024
		Global closed-canopy coconut palm	20m	2020	Descals et al. 2023
	Short vegetation, bare	Global Pasture Watch (GPW) Cultivated Grasslands	30m	2020	Parente et al. 2024
	Cropland	USGS Global Cropland Extent Product at 30m Resolution (GCEP30)	30m	2015	Thenkabail et al. 2021
		ESA WorldCereal Active Cropland	10m	2020	Van Tricht et al. 2023
	Built-Up	Mining Polygons	Vector	2019, 2020	Maus et al. 2022, Tang et al. 2024, Dethier et al. 2023
Natural	Forests	Intact Forest Landscapes (IFL)	Vector	2020	Potapov et al. 2017
	Mangroves	Global Mangrove Watch (GMW), version 3.0	0.8 arc seconds	2020	Bunting et al. 2022

## Forests

The UMD forest class includes all tree cover greater than or equal to 5 meters in height, regardless of whether it is planted or natural. Tree cover is a convenient metric for monitoring forests because it is easily measurable from space, but cannot be used to assess natural forests on its own. While no globally consistent planted or natural forest dataset exists, they can be delineated through the use of multiple ancillary datasets. Here we applied three additional datasets to identify non-natural and natural forest in the UMD forest class.

### *The Spatial Database of Planted Trees (SDPT), version 2.0*

The SDPT is a global dataset of tree crops – defined as stands of perennial trees for agricultural purposes – and planted forests – defined as stands of planted trees (other than tree crops) grown for wood and wood fiber production or for ecosystem protection against wind and soil erosion (Richter et al. 2024). This is a vector dataset of compiled and harmonized national or regional maps from a variety of sources, including national governments, nongovernmental organizations, independent researchers, or a combination of sources. As such, the resolution, methods, year, and accuracy of input data vary by source (see Appendix D for a full list of the input data for the SDPT v2, as well as their native resolution and year).

Tree crops and tree plantations do not meet the AFi definition of a natural forest. Although “planted forest” as defined in the SDPT may in some instances meet the AFi definition of natural forest (e.g. if natural species composition, structure, and function is maintained), the SDPT specifically includes plantations that were likely to be intensively managed and excludes areas of semi-natural forest with natural regeneration. Therefore, we consider “planted forests” in this dataset to represent a reasonable proxy of “tree plantations” as defined by AFi. Version 2.0 of this dataset is used to classify forests as non-natural (Richter et al. 2024).

### *Global closed-canopy coconut palm*

To supplement the SDPT v2, we incorporated a global map of closed-canopy coconut palm (Descals et al. 2023). These data were produced using a semantic segmentation model to classify Sentinel-1 and Sentinel-2 annual composites for 2020 within regions that are suitable for the cultivation of coconut palms based on a bioclimatic analysis. Closed-canopy coconut palm was defined as including more than four coconut palms within a 20 meter pixel, with coconut palms fully covering the ground (e.g. reaching full canopy closure). Because tree crops do not meet the AFi definition of a natural forest, we used these data to classify forests as non-natural.

### *Intact Forest Landscapes (IFL)*

Intact Forest Landscapes are defined as mosaics of forests and naturally treeless ecosystems within the zone of current forest extent that show no signs of significant human activity or habitat fragmentation and are large enough to maintain all native biodiversity (Potapov et al. 2017). These data map the extent of Intact Forest Landscapes globally in 2020. Forests within Intact Forest Landscapes are likely to meet the AFi definition of natural, as they show no signs of significant human activity. Therefore, we used the IFL extent in 2020 to apply an additional precautionary measure to ensure forests that fall within these boundaries are classified as natural.

**Short Vegetation and Bare**

A key challenge in mapping natural short vegetation and bare ground is distinguishing it from pasture or other grasslands that have been significantly transformed or degraded by livestock grazing. To help delineate these areas, we used ‘cultivated grassland’ data from the Land & Carbon Lab’s Global Pasture Watch (GPW). Cultivated grasslands are areas where grasses and other forage have been intentionally planted and managed as well as areas of heavy management for human-directed uses such as livestock grazing. These landscapes often exhibit a high level of human intervention through practices such as regular seeding, fertilization, controlled grazing, and sometimes irrigation, aimed at enhancing productivity and maintaining the desired vegetation cover. The presence of livestock-related infrastructure such as fencing is typical, reflecting the managed nature of these areas (Parente et al. 2024). The GPW data were created by predicting global grassland areas separately in two classes: cultivated and natural/semi-natural. Predictive models to map the grass classes, separately, were fitted using a random forest classifier, resulting in global probabilities maps at 30 meter spatial resolution. A resulting combined dominant class map was created.

**Wetlands**

Both the forest and short vegetation classes have areas labeled as wetland or peat. The wetland label comes from overlaying the UMD wetland class and may include inland, coastal, saline, freshwater, and brackish wetlands. Areas with mangroves are labeled as mangrove instead of forest. These are wetland types that are of high-interest to map users because of their high potential for carbon storage. Mangroves are designated as a unique class using data produced by Global Mangrove Watch (GMW) on mangrove extent for the year 2020 (Bunting et al. 2022). The GMW data was produced using L-band Synthetic Aperture Radar (SAR) global mosaic datasets from the Japan Aerospace Exploration Agency (JAXA) for 11 epochs from 1996 to 2020 to develop a time-series of global mangrove extent and change. Although the ESA WorldCover data include a separate class of mangrove forests, the GMW data were selected because of their high accuracy scores (86% producer’s accuracy and 89% user’s accuracy) (Bunting et al. 2022). Peatland extent was included by using a map developed by researchers at WRI modeling forest carbon emissions and removals (Harris et al. 2021). This map (WRI Peat) is a 30 meter resolution composite of 5 peatland maps that were either converted from vector data to raster data, or were resampled from coarser resolution raster data. This composite includes 3 datasets with regional coverage and 2 with global coverage. This peat map was overlaid with the forests and short vegetation class to delineate peat forests and peat short vegetation.

Table 6. Input data used to create the composite peat map.

Coverage	Native Resolution	Source
Indonesia and Malaysia	Vector	Miettinen et al. 2015
Congo Basin	50m	Crezee et al. 2022
Lowland Peruvian Amazon	50m	Hastie et al. 2022
Land area below 40 degrees north	250m	Gumbrić et al. 2017
Land area above 40 degrees north	Vector	Xu et al. 2018

## Cropland

To supplement the UMD Land Cover cropland class, we used two additional global cropland datasets.

### *USGS Global Cropland Extent Product (GCEP30)*

The USGS Global Cropland Extent Product at 30 meter resolution (GCEP30) for the year 2015 (Thenkabail et al. 2021) were developed through the classification of Landsat imagery using machine learning algorithms trained for 74 agroecological zones and compiled into one global map (Thenkabail et al. 2021). In this dataset, cropland includes the following: cropland cultivated one or more times throughout a 12-month period, cropland that is left fallow but is equipped for agriculture, and cropland that is permanently cropped with plantations (such as vineyards, orchards, coffee, tea, etc.). In Argentina, Australia, Brazil, Kazakhstan, and New Zealand, the data also include managed pasture, as is likely in much of South and Central America. Upon visual inspection in Mexico and Nigeria, we found the GCEP30 data included large areas of agricultural lands (possibly managed pasture, mixed crop/pasture, fallow fields or abandoned cropland) that were not included in the UMD cropland extent. Because the UMD data limit the fallow period to four years, we incorporated the GCEP30 data to include cropland that may experience longer fallow periods, as well as managed pasture in the regions where it is included.

### *ESA WorldCereal Active Croplands*

The ESA WorldCereal consortium created a series of global mapping products for temporary cereal grains including seasonality and management systems for the year 2021 using Sentinel-1, Sentinel-2, and Landsat 8 (Van Tricht et al. 2023). They mapped active croplands regardless of crop type before classifying cereals. They define temporary crops as crops “with a less-than-1-year growing cycle which must be newly sown or planted for further production after harvest,” however they do also include sugar cane, asparagus, and cassava (Van Tricht et al. 2023). The WorldCereal active cropland layer maps temporary crops that have an entire growing cycle in a specific growing season based on existing crop calendars. For the 2020 Natural Lands Map, we used active cropland areas that have growing seasons that began in 2020 and excluded growing seasons that began in 2021 to ensure we were not mapping any new cropland that was established in 2021.

## Built-up

To supplement the UMD Land Cover built-up class, we combined three datasets delineating mining boundaries into a single layer using a spatial union: IIASA Global Scale Mining Polygons (Maus et al. 2022) for the year 2019, Global Mining Footprint (Tang et al. 2023) for the year 2020, and Global Alluvial Mining (Dethier et al. 2023) for the year 2020. All three datasets were created through manual delineation of mining areas using satellite imagery. Some mining areas contain a mix of mining activities and natural lands such as tree patches, so only the areas within mining boundaries that are bare and water land covers were used to identify non-natural areas.

### 2.2.3. REGIONAL DATA

We evaluated regional data using our criteria for inclusion described in Table 3. While not all regional data have accuracy assessments, all contain more detailed or regionally specific classes. These data were harmonized with our map classes and incorporated into the map, replacing our global data where available. In cases where regional data included only one or

a few classes relevant for our natural lands map (e.g. cocoa, cropland, primary forest, and natural grassland/shrubland maps), these data were used to supplement or replace the relevant map class.

Table 7. Summary of regional datasets incorporated into the map.

Region	Dataset Name	Year	Reference	Resolution	Classes incorporated in natural lands map
Brazil	MapBiomias Brazil Collection 9.0	2020	Souza et al. 2020	30m	All classes
Amazon	MapBiomias Amazonia Collection 6.0	2020	MapBiomias Amazonia 2024	30m	All classes
Indonesia	MapBiomias Indonesia Collection 2.0	2020	MapBiomias Indonesia 2024	30m	All classes
Peru	MapBiomias Peru Collection 2.0	2020	MapBiomias Peru 2024	30m	All classes
Bolivia	MapBiomias Bolivia Collection 2.0	2020	MapBiomias Bolivia 2024	30m	All classes
Colombia	MapBiomias Colombia Collection 2.0	2020	MapBiomias Colombia 2024	30m	All classes
Venezuela	MapBiomias Venezuela Collection 2.0	2020	MapBiomias Venezuela 2024	30m	All classes
Uruguay	MapBiomias Uruguay Collection 2.0	2020	Baeza et al. 2022	30m	All classes
Ecuador	MapBiomias Ecuador Collection 2.0	2020	MapBiomias Ecuador 2024	30m	All classes
Paraguay	MapBiomias Paraguay Collection 1.0	2020	MapBiomias Paraguay 2024	30m	All classes
Chile	MapBiomias Chile Collection 1.0	2020	MapBiomias Chile 2024	30m	All classes
Argentina	MapBiomias Argentina Collection 1.0	2020	Mapbiomas Argentina 2024	30m	All classes
South Africa	South Africa National Land Cover 2020	2020	Department of Forestry, Fisheries, and the	20m	All classes



			Environment, South Africa		
Côte d'Ivoire and Ghana	ETH/EcoVision Cocoa Map	2019-2021	Kalischek et al. 2023	10m	Single class
New Zealand	LUCAS NZ Land Use Map	2020	Ministry for the Environment	Vector	All classes
Europe	CORINE Land Cover	2018	Copernicus Land Monitoring Service	100m	Select classes: Natural grassland, moors and heathland, sclerophyllous vegetation
Europe	European Primary Forest Database (EPFD) version 2.0	Varies	Sabatini et al. 2021	Vector	Single class
United States	U.S. Geological Survey National Land Cover Database (NLCD)	2020	USGS 2024	30m	Select classes: Open space, low intensity developed, medium intensity developed, high intensity developed, pasture/hay, cultivated crops
Africa	Digital Earth Africa (DEA) cropland extent	2019	Digital Earth Africa	10m	Single class

MapBiomass Land Cover and Land Use products for Brazil, the Amazon, Peru, Bolivia, Colombia, Venezuela, Uruguay, Ecuador, Paraguay, Chile, Argentina and Indonesia are 30 meter resolution maps that contain detailed land cover/land use classes for natural forest and non-forest ecosystems, as well as agriculture, water, and non-vegetated areas. We used the year 2020 Land Cover and Land Use product for all MapBiomass collections.

The South Africa National Land Cover map for 2020 was produced using multi-seasonal 20 meter resolution Sentinel-2 satellite imagery. This map contains 73 classes that delineate natural and non-natural land covers.

The Ghana and Côte d'Ivoire cocoa map for 2019-2021 was produced using Sentinel-2 satellite imagery at a 10 meter resolution. The map delineates land under cocoa cultivation, including shade grown cocoa.

The Land Use and Carbon Analysis System (LUCAS) New Zealand Land Use data for 2020 are based on Sentinel-2 satellite imagery acquired in the summer of 2020/2021. There are 30 distinct land use classes, including classes for natural forests and natural grasslands.

The CORINE Land Cover dataset for 2018 is a complete land cover map over the participating countries of the European Environment Agency at a 100 meter resolution. We included CORINE data to improve the delineation of natural grasslands in Europe. We used three shrub classes: natural grassland, moors and heathland, sclerophyllous vegetation, as natural short vegetation.



The European Primary Forests Database (EPFD) defines primary forests as forests where the signs of human impacts, if any, are strongly blurred due to decades without forest management (Sabatini et al. 2021). These data combine and harmonize 48 datasets of primary forests in 33 countries in Europe, and were used to aid in the delineation of natural forests in Europe. While these data include both polygons and point features, only polygons were used for the natural lands map. Due to the variety of data sources used in these data, data quality, accuracy, and completeness vary.

The United States Geological Survey Annual National Land Cover Database (NLCD) for 1985–2023 maps land cover and change across the contiguous United States based on Landsat composites. We used classes in the 2020 map that help delineate non-natural lands: developed (open space, low intensity, medium intensity, and high intensity), pasture/hay, and cultivated crops, which were reclassified to built-up, non-natural short vegetation, and cropland respectively.

The Digital Earth Africa cropland extent map for 2019 maps cropland over the entire African continent. Cropland is defined as a piece of land that is sowed/planted and is harvestable at least once within 12 months after the sowing/planting date. Non-planted grazing lands and perennial crops are excluded. The map was produced using Sentinel-2 satellite images for 2019, and each region was mapped separately and then combined. We used the ‘filtered’ product, which applies an image segmentation algorithm (Clewley et al. 2014) to the cropland extent mask in order to merge segments smaller than 1 hectare with neighboring segments.




Figure 1. Extent of regional data included in the natural lands map.

## 2.3. Methods

### 2.3.1. METHODS FOR CREATING AND COMBINING MAP CLASSES FOR GLOBAL DATA

To create the global 2020 natural lands map, we combined our input data through a series of overlays and decision rules (Figure 2). The map has a hierarchical legend, with level 1 distinguishing two classes– natural and non-natural– and level 2 distinguishing various land cover classes within the natural and non-natural classifications. The map includes both natural and non-natural forests, short vegetation, water, bare ground, wetland forests, peat forests, wetland short vegetation, and peat short vegetation. Natural classes also include mangroves and permanent snow/ice. Non-natural classes also include built-up and cropland.

Table 8. Final map classes, values, and description

Level 1	Level 2	Class Value	Description
Natural	Forest	2	Tree cover greater than 5 meters in height and more than 0.5 hectares, excluding planted forests grown for wood or wood fiber production or perennial tree crops. Height or minimum mapping thresholds may vary based on local definitions.
	Short vegetation	3	Areas of land with vegetation shorter than 5 meters (including areas of land dominated by grass or shrubs), but excluding areas with cultivated grasslands, cropland, or tree crops.
	Water	4	Surface water present 20% or more of the year, where water is the dominant class.
	Mangroves	5	Areas dominated by mangrove forests.
	Bare	6	Areas with exposed rock, soil, or sand with less than 10% vegetated cover but excluding areas with high densities of ruminant livestock, cropland, or tree crops.
	Snow/Ice	7	Land covered by glaciers or snow remaining during the entire year.
	Wetland forest	8	Natural forests with saturated soil that can be inundated by water either seasonally or permanently.
	Peat forest	9	Natural wetland forests that have accumulated peat.
	Wetland short vegetation	10	Natural short vegetation with saturated soil that can be inundated by water either seasonally or permanently.
Peat short vegetation	11	Natural wetland short vegetation that have accumulated peat.	
Non-natural	Cropland	12	Land used to produce annual and perennial crops. This may include herbaceous crops, land that is permanently cropped with plantation crops (such as vineyards, coffee, tea), some tree crops (such as orchards), and some pasture or mixed pasture/crop systems. It includes land that is left fallow but equipped for agriculture.

Built-up	13	Man-made land surfaces associated with infrastructure, commercial, residential uses, and mining.
Tree cover	14	Perennial tree crops (including rubber, oil palm, cocoa, orchards, etc.) and planted forests grown for wood or wood fiber production. This may include both intensively managed forest plantations for timber with a short rotation time, or managed forests with signs that the forest has been planted with a long rotation time (greater than 15 years).
Short vegetation	15	Pasture, tree or plantation crops, or other areas with vegetation shorter than 5 meters that are cultivated grasslands.
Water	16	May include aquaculture, artificial dams, or other artificial areas with surface water.
Wetland tree cover	17	Non-natural tree cover with saturated soil that can be inundated by water either seasonally or permanently.
Peat tree cover	18	Non-natural wetland tree cover that have accumulated peat.
Wetland short vegetation	19	Non-natural short vegetation with saturated soil that can be inundated by water either seasonally or permanently.
Peat short vegetation	20	Non-natural wetland short vegetation that have accumulated peat.
Bare	21	Areas with exposed rock, soil, or sand with less than 10% vegetated cover but with a high density of ruminant livestock.




Figure 2. Process used to create the natural lands map. The diagram shows the order in which we conducted each step; the details of each step in the process can be found in the Methods section. As an example, the first line in the diagram shows the following process for separating natural forest from non-natural tree cover: the forest class from the UMD Global Land Use and Land Cover Change dataset was designated as natural forest. Any forest areas within the IFL 2020 layer and EPFD layer were kept as natural forest. Any forest area outside of those two layers which overlap with SDPT v2, ETH Cocoa data, or the Global Coconut data were reclassified as non-natural tree cover. A minimum mapping unit (MMU) of 0.5 ha was then applied to the natural class; any forest patches that did not meet the 0.5 ha MMU were classified as non-natural tree cover.

We applied a conservative approach in mapping non-natural lands, meaning that decisions were made with the aim to be precautionary in assigning a non-natural classification to an ecosystem. As a result, our final data may overestimate the area of natural lands in some regions. Due to the lower resolution and variation in accuracy of some of our input data, we used additional data where available to apply additional conditions before removing non-natural classes as an added precautionary step. Because our map may overestimate natural lands, we recommend that this map only be used for applications that align with the

definitions of ‘natural’ used in the map, such as setting a corporate “no conversion of natural ecosystems” target in SBTN Land, and not used to quantify the area of natural or non-natural lands.

We adopted this conservative approach because the risk of underestimating natural lands is greater than the risk of overestimating natural lands for the protection of nature. If natural lands were underestimated, natural areas not included in the map may be at higher risk of conversion because they are not designated as “off limits” for conversion. Particularly considering the data limitations for certain land cover types – including grasslands and pasture – we considered it more appropriate to adopt a precautionary approach so that potential natural lands are rarely omitted from the map. This approach can be beneficial to companies and other entities, too, since a conservative approach makes it less likely to unknowingly convert natural lands. However, the overestimation of natural lands is unevenly distributed, with more natural lands in areas with less data to distinguish between natural and non-natural.

All processing steps and analyses were conducted in Google Earth Engine (Gorelick et al. 2017).

### **Pre-processing**

All data were converted to raster format and resampled to 30 meter resolution to match the resolution of the UMD Land Cover data, which were used as the base layer for the majority of our land cover classes. The data that had higher resolutions were resampled by using the mode pixel value within each 30 meter pixel. We resampled the lower resolution data to fit the projection and resolution of the 30 meter data.

### **Forests**

To delineate forests, we used the tree height layer from the UMD Land Cover data, which define tree cover as all woody vegetation, and filtered the data to tree cover greater than or equal to 5 meters in height. All forest within the UMD forest extent was assumed to be natural unless re-classified as non-natural. Because these data include both natural forests and non-natural tree cover, such as wood fiber plantations or tree crops, we relied on the SDPT version 2.0 (Richter et al. 2024) and the global map of closed-canopy coconut palm (Descals et al. 2023) to target non-natural tree cover.

To remove tree crops and planted forest from the natural forest class, we overlaid the SDPT v2 and closed-canopy coconut palm map (combined via union) with the UMD forest class and reclassified any areas of overlap as non-natural tree cover. However, due to regional variations in the resolution and accuracy of the SDPT v2.0 source data, which limits the precision and accuracy by which non-natural tree cover can be delineated in certain cases, we used data on the extent of intact forest landscapes from the global IFL data in 2020 to apply additional conditions before reclassifying forest as non-natural. These data, which map forests with no signs of human intervention, are likely to represent areas containing natural forests. In cases where the SDPT overlapped with areas designated as intact forest landscapes, the IFL data were given priority and the forest remained classified as natural.

Finally, to align with the AFi definition of forest, we applied a minimum mapping unit of 0.5 hectares to the natural forest class. 0.5 hectares is approximately 5.6 pixels, meaning that if there is a patch of natural forest smaller than 0.5 hectares (patches can be connected by any side or corner of the pixel), it is labeled non-natural tree cover, unless another class with higher priority re-labels it another class when the data are all compiled.

## Short vegetation and bare

With limited global data available regarding natural and non-natural short vegetation (including grasslands and shrublands) and bare ground, we used cultivated grasslands as a proxy for pasture. We used unions to combine the ESA WorldCover grassland, shrubland, and herbaceous wetland classes as the extent of short vegetation, and the ESA WorldCover barren/sparse vegetation class with the UMD bare class as the extent of bare ground. All short vegetation and bare ground were assumed to be natural unless re-classified as non-natural. We used the GPW 2020 dominant map, which combined pixels that have natural/semi-natural grasslands probabilities over 43% or cultivated probabilities over 32% into one map, where the natural/semi-natural has priority. From the dominant map, we used the area identified as cultivated grasslands. We then classified areas identified as both cultivated grasslands (GPW) and short vegetation or bare (ESA WorldCover and UMD) as non-natural. Any areas of short vegetation or bare that overlapped with the closed-canopy coconut palm data and plantations from the SDPT (after removing any overlap with IFL) were also labeled as non-natural, since orchards and other tree crops may be shorter than 5 meters in height and therefore classified as short vegetation.

## Water

Water was mapped with the surface water dynamics layer from the UMD Land Cover data, which includes all water present 20% of the year or more in 2020. Because data distinguishing natural from non-natural water was not available or not deemed appropriate, at the global level, water bodies were classified as natural by default.

## Built-up

The UMD Land Cover built-up class was used as the primary extent of built-up area. We also assigned the built-up class in areas within the combined global mining polygons that were labeled as water or bare as defined above. This combination covered most mining activities, while preserving areas of natural forest or short vegetation within the polygons.

## Combining map classes

For the cropland class, we combined the UMD cropland class with the GCEP30 cropland data and ESA WorldCereal active cropland data by taking the union of all three datasets. All the final class layers as described above were compiled with the following priority from highest to lowest: built-up, cropland, non-natural tree cover, non-natural short vegetation, non-natural bare ground, mangroves, natural forest, natural short vegetation, natural water, bare ground, snow/ice, and non-natural water (which is present in regional data). In cases where any classes overlap, the higher priority class takes precedence. Since we assume all land to be natural unless re-classified as non-natural, and we are generally conservative in assigning a non-natural class, we prioritized non-natural land uses when compiling the map. We filled in any remaining empty pixels caused by using land cover data from different sources as natural short vegetation.

## 2.3.2. METHODS FOR INCORPORATING REGIONAL DATA

In creating the natural lands map, we aimed to include regional or country-level data where available and appropriate. Therefore, where quality regional data are available that meet our criteria, we harmonized these data to our global classes and used them in place of our global data. For this version of the map, we incorporated MapBiomas data for Brazil, the Amazon, Bolivia, Colombia, Venezuela, Uruguay, Ecuador, Paraguay, Chile, and Argentina in South



America, and Indonesia. For overlapping MapBiomass datasets in South America, the following layering priority was used, from highest to lowest: Brazil, Ecuador, Venezuela, Colombia, Bolivia, Peru, Chile, Argentina, Paraguay, Uruguay, and the Amazon. We also incorporated the South Africa National Land Cover Map, and the New Zealand LUCAS Land Use map, the developed and agriculture classes from the US National Land Cover Database and the grassland classes from the European CORINE Land Cover Data. We also incorporated data that corresponded to a single class in the natural lands map, including the European Primary Forest Database, the ETH/EcoVision Cocoa Map for Côte d'Ivoire and Ghana, and the Digital Earth Africa (DEA) cropland extent map (see Table 7).

### **Regional data using single or select classes**

For regional data with only one or few classes that correspond to the natural lands map (e.g. the ETH/EcoVision Cocoa map, the DEA cropland extent map, the European Primary Forest Database, USGS National Land Cover Database, and the CORINE Land Cover data), these data were reclassified to the corresponding map class and incorporated into the map at the same stage as the global data and following the same layering priority described in section 2.3.1. Areas mapped as cocoa by the ETH/EcoVision Cocoa map were classified as non-natural tree cover, and then layered into the map with the built-up and cropland classes given priority in any areas of overlap. All areas of overlap between short vegetation and CORINE natural shrub classes were labeled as natural short vegetation, while all areas of overlap between the forest class and the European Primary Forest Database were labeled as natural forests. The USGS National Land Cover Database developed (open space, low-, medium-, and high-intensity), pasture/hay, and cultivated crops were reclassified to built-up, non-natural short vegetation, and cropland, respectively, and replaced the global data in the map. DEA cropland was layered into the map following the same layering priority as the global cropland layers.

### **Regional data using all classes**

For regional data where wall-to-wall land cover maps were incorporated (MapBiomass collections, South Africa National Land Cover Map, and New Zealand LUCAS Land Use map), we reclassified each dataset according to the classes used in our global map (see Appendix A, B, and C) and replaced the global data with the harmonized regional data. Where possible, we applied a direct reclassification to convert each class in the regional dataset to the class that was most closely aligned in our global dataset. For classes that contained both natural forest and natural non-forest vegetation according to the AFI/FAO definitions (such as savannas, woodlands, or other mixed classes), we overlaid the UMD tree height data (greater than or equal to 5 meters) to delineate natural forest versus natural short vegetation. As with the forest data globally, we applied a minimum mapping unit of 0.5 hectares and classified any patches that did not meet this threshold as natural short vegetation. We applied this method to the savanna class in the MapBiomass Brazil, Amazon, Peru, Venezuela, Ecuador, and Paraguay collections; as well as the sparse woodlands class in the MapBiomass Chaco collection. It was also applied to the contiguous low forest & thicket, dense forest & woodland, and open woodland classes in the South Africa National Land Cover map.

We determined that in several regional datasets, the definitions of certain classes were too broad to be directly reclassified as a single class in our map, and in some cases encompassed both natural and non-natural areas. In these cases, we used the UMD Land Cover data to assist in reclassifying the broad class into multiple classes that are harmonized with our global map. For example, the MapBiomass Uruguay, Argentina, and Paraguay datasets have a single class for non-vegetated areas, which can include both non-natural built-up areas or exposed soil in cropland, or natural areas of bare land, rock or sand. Likewise, the 'other

non-vegetated areas' class in MapBiomias Peru, Indonesia, and Chile includes both natural and non-natural areas. In these cases, we reclassified this non-vegetated class as natural bare land unless 1) it overlapped with the UMD built-up class, in which case we reclassified it as built-up; or 2) it overlapped with the UMD cropland class, in which case we reclassified it as cropland. In Brazil, the Amazon, Bolivia, Colombia, Ecuador, and Venezuela collections the 'other non-vegetated areas' or 'other non-vegetated anthropic areas' classes were defined as all non-natural land covers. We re-classified these areas as non-natural bare land unless it overlapped with the UMD built-up or cropland classes in order to have a specific class. Similarly, MapBiomias Indonesia was the only MapBiomias collection without a wetlands class. In this case, we reclassified the 'other non-forest formations' class as natural short vegetation unless it overlapped with the UMD wetland short vegetation class, in which case we reclassified it as natural wetland short vegetation. This was also done with the New Zealand LUCAS Land Use data to separate the permanent snow and ice from the rest of the bare class.

### 2.3.3. METHODS FOR CREATING THE FINAL MAP

After replacing our global data with regional data where applicable, the forest and short vegetation classes (both natural and non-natural) were overlaid with the WRI Peat data and assigned a peat label where applicable. The final result is a land cover map with forest, short vegetation, mangroves, water, bare, snow/ice, built-up, and cropland, where the forest, short vegetation, bare, and water have both natural and non-natural classes, and forest and short vegetation are labeled as dry, wetland, or peat. For the use of the SBTN no conversion target, the level 1 categories in the map can be used to create a binary image: natural and non-natural.

### 2.3.4. VALIDATION METHODS

International Institute for Applied Systems Analysis (IIASA) conducted an independent accuracy assessment of the natural lands map during the creation of the beta version (v0.3). They created a validation data set of natural and non-natural classes using a simple random sample of 4,943 points globally. Each of the 4,943 points were validated by two IIASA experts with visual inspection of very high resolution imagery in a Geo-Wiki web-application created with various supplementary data, including Google imagery, Microsoft Bing imagery, Esri images, NDVI time series, Sentinel 2-time series, etc. The validation team followed the operational definitions used in the map for the natural/non-natural classes to guide decisions applied for the labeling of validation data. To account for geolocation errors in both the original map as well as the underlying very high-resolution imagery, additional neighboring pixels around the central pixel were classified totaling 9 sub-pixels. Both the majority class and middle pixel class were used in the validation (Xu et al. 2024). Any disagreements in the classification of the two experts were revised by a third expert. In some locations, very high-resolution imagery was either not available or not frequently available, so it could be difficult to determine the class of a given validation point, and a label of "not sure" was given. Points with a "not sure" label were not included in the results of the accuracy assessment. The overall and per class accuracies were derived from confusion matrices at a global scale. Using the validation points labeled natural and non-natural, we retrieved the values from the V1.1 binary SBTN Natural Lands Map and calculated a confusion matrix with User's, Producer's, and Overall accuracies.

Because of the specific interest in the natural forest class for deforestation monitoring, IIASA created a validation dataset to assess the accuracy of the natural forest class. The validation sample was a subset of the original Natural Lands Map validation dataset of all points that fell within potential natural forest areas, which were delineated through the combination of three forest datasets: World Cover 2020 tree cover, Meta/WRI High



Resolution 1m Global Canopy Height for 2020 (Tolan et al. 2024), and natural forests (including the wetland, peat, and mangrove natural forest subclasses) from version 1 of the SBTN Natural Lands Map. All maps were aggregated to 1 km by classifying a 1 km cell as potential forest area if at least one pixel from any input datasets fell within the cell. This resulted in a sample of 3,242 validation points. Each point was interpreted by two experts using the natural forest definition in the SBTN Natural Lands Map. Any disagreements were revised by a third expert.

# 3. Results

## 3.1. Map Results

The natural lands map shows that large blocks of natural land still exist across most regions of the world. Figure 3 shows the map of natural (green) and non-natural (gray) areas. The majority of non-natural land is built-up areas, cropland, and pastureland. There are also large contiguous regions of non-natural tree cover. As expected with our conservative approach in designating non-natural lands, visual inspection showed that some areas of non-natural tree cover were classified as natural. However, our visual inspection indicated that there were not many obvious places where natural forests were classified as non-natural, indicating that our conservative definition of non-natural tree cover produced the intended result.




Figure 3. Global map of natural lands. Note: There is no data on the glaciers of Greenland.

In Figure 4 we highlight six regions around the world. The natural lands are broken out by land cover and the non-natural lands were combined into one class and shown in gray. Figure 3a) shows the Cerrado in Brazil, which is a mix of short vegetation and low density forests, and non-natural areas, which are largely agricultural croplands, pasturelands and plantations. The north-west corner is the Amazon rainforest and the north-east corner is the arid Caatinga savanna. Figure 3b) focuses on Colombia and depicts natural forests in the Amazon rainforest and the Andean region. There is natural short vegetation in the Orinoquia region in the northeast; however, there are also non-natural lands occupied by pasture. Figure 3c) covers some of western/southern Europe centered on the Alps. The natural lands map classifies many areas of this region as non-natural, with the exception of some natural forests in Germany, France, Switzerland, and Italy, and the mountainous areas of the Alps, which are mostly bare land, permanent snow/ice, short vegetation, and forest. Figure 3d) shows natural forests and short vegetation in West Africa, with built-up areas, cropland, and tree plantations - largely cocoa - categorized as non-natural land. Most of Figure 3e) in the Congo basin is natural forest, with peat forest in the middle and some natural short vegetation to the south. There are non-natural areas in the northern part of the image that are mostly cropland with some built-up areas. Finally, Figure 3f) shows Peninsular Malaysia's natural forests and the non-natural area, which is mostly oil palm plantations.




Figure 4. Natural land covers in a) Brazil; b) Colombia; c) western/southern Europe; d) Côte d'Ivoire; e) the Congo basin; f) Peninsular Malaysia.



Upon visual inspection, areas where we replaced the global data with regional data improved significantly. Figures 5a) and 5b) depict part of the Manawatu-Whanganui region of the north island of New Zealand. Figure 5a) shows the version with only global data, which maps mostly non-natural short vegetation in the eastern half of this region, whereas 5b) (after incorporation of LUCAS data) designates these areas identified as low producing grasslands in the LUCAS data as natural short vegetation. Figures 5c) and 5d) are centered on an area of southern Côte d'Ivoire and Ghana that have many cocoa plantations. Figure 5c) shows the version with only global data, which shows mostly natural forests in the area, whereas 5d) (after incorporation of ETH/EcoVision Cocoa Map) designates lots of areas as non-natural. The cocoa data helps identify plantations in this area that were not captured by the Spatial Database of Planted Trees. Bringing in regional data improved the natural lands map through the use of datasets with higher accuracy that were produced to suit the regional context and based on local knowledge of these landscapes.




Figure 5. Comparison of area in a) New Zealand with global data vs. b) New Zealand with the incorporation of LUCAS data; and the comparison of area in c) Côte d'Ivoire and Ghana with global data vs. d) Côte d'Ivoire and Ghana with the incorporation of ETH/EcoVision Cocoa Map data.

### 3.2. Validation and accuracy

The natural and non-natural binary map as validated by IIASA showed a 92.1% overall accuracy. Three percent of the validation points were classified as “not sure” because of a lack of high resolution imagery available from 2020, and were removed before calculating the accuracy. The natural class had a 96.3% User’s accuracy and 93.8% Producer’s accuracy. These results show that the map mis-classifies 6% of the natural points as non-natural, and 15% of the non-natural points as natural. This result is expected, as our mapping approach was precautionary in assigning a non-natural label.

Table 9. Confusion matrix showing agreement between validation dataset and natural lands map.

		Reference			
		Natural	Non-natural	Total	User’s Accuracy
Map	Natural	3590	139	3729	96.3%
	Non-natural	236	772	1008	76.6%
	Total	3826	911	4737	
	Producer’s Accuracy	93.8%	84.7%		92.1%

The natural forests class had a User’s and Producer’s accuracy of 87.2% and 85.0%, respectively. When the natural forest class was compared against all other classes, there was an overall accuracy of 88.8%.

Table 10. Confusion matrix showing agreement between validation dataset and natural forest class.

		Reference			
		Natural Forest	Other	Total	User’s Accuracy
Map	Natural Forest	1129	165	1294	87.2%
	Other	199	1746	1945	89.8%
	Total	1328	1911	3239	
	Producer’s Accuracy	85.0%	91.4%		88.8%

### 3.3. Comparison with existing data

Gosling et al. (2020) produced a global map at 1 km resolution of natural and modified habitat for use in investment screening as part of Performance Standard 6 (PS6) of the International Finance Corporation (IFC) by combining eleven data layers. IFC PS6 defines natural habitats as “areas composed of viable assemblages of plant and/or animal species of largely native origin, and/or where human activity has not essentially modified an area's primary ecological functions and species composition” and modified habitats as “areas that may contain a large proportion of plant and/or animal species of non-native origin, and/or where human activity has substantially modified an area's primary ecological functions and species composition”. Gosling et al. (2020) use these definitions to classify and combine input data, relying on human pressure as a proxy for the loss of ecological function and species composition. Aside from differences in input data used, Gosling et al.'s approach differs in a few ways: 1) they use only global data, whereas our approach incorporates regional or regional data where available; 2) they overlay data representing natural and modified categories, and fill in remaining area (37.5% of global land area) using a categorized Human Footprint Layer (Venter et al. 2016); whereas our approach starts with land cover classes with global coverage and uses supplementary data to remove non-natural areas.

Gosling et al. used four categories in their map, representing a gradient between natural and modified: 1) likely modified, 2) potential modified, 3) potential natural, and 4) likely natural. We reclassified these four categories into two categories for better comparison with our map. Likely modified and potential modified were reclassified as “non-natural”, and likely natural and potential natural were reclassified as “natural”. We resampled our natural lands map to match the 1 km resolution of Gosling et al. map, taking the mode value of our binary layer.

Overall, the maps had high agreement: 59% of area was classified as natural by both maps and 20% of area was classified as non-natural by both maps, resulting in 79% overall agreement. However, a larger percentage of our map was classified as natural: 19% of the area classified as natural by our map was classified as non-natural by Gosling et al. Meanwhile, only 3% of the area classified as non-natural by our map was classified as natural by Gosling et al. Although Gosling et al. similarly take a precautionary approach by prioritizing natural categories when there is disagreement between the input datasets used, they incorporate the Human Footprint Layer, which uses data on population density and proximity to roads, variables which were not considered in our natural lands map. This is likely a key source of disagreement between the two maps.

# 4. Limitations

Users of the map should be cautious of its limitations, and should take additional steps, such as validation with high resolution imagery or ground-truthing or use of additional data, to supplement the use of the map, especially for local applications. The natural lands map includes a number of important data limitations:

1. **Definitional inconsistencies:** The dataset definitions do not always match the definitions outlined by AFi. AFi provides robust definitions of natural forests; however, other natural ecosystems, including those dominated by short vegetation, wetlands, water, and snow and ice lack the same level of distinction, and as a result we relied on definitions derived from the data used to create the map. There are also definitional inconsistencies across various data sources used to create the natural lands map, which is a tradeoff for including regional data when possible (see below for more detail). For example, the SDPT data, used to exclude tree plantations from natural forests, include dozens of local sources. While most capture short rotation plantations and tree crops, they also include mixed use areas dominated by tree plantations. Similarly, the MapBiomass data do not always have a height threshold used to define forests, which may create inconsistencies with the delineation of forests within regions in which Mapbiomas data was used, versus those which relied exclusively on the UMD forest extent

Additionally, it should be noted that not all livestock grazing areas are captured within the cultivated grassland data, and many semi-natural areas with livestock grazing remain in the natural class. The impact that livestock grazing can have in these areas falls along a continuum from minimal to severe and depends not only on the density of livestock but also the climate, soil, and evolutionary history of grazing in that area. AFi does not have a specific definition for the natural short vegetation or grasslands, and because GPW data is the only relevant global dataset at this time, the operational definition of natural short vegetation was based on what was captured in the cultivated class.

2. **Temporal inconsistencies:** While the map is as close to the year 2020 as possible, some data are from earlier time periods. For example, the USGS National Land Cover Database is from 2019 and the USGS Global Cropland Extent Product is from 2015. Likewise, some source datasets for the SDPT and EPFD are from earlier time periods. See section 2.2 and Appendix D for more details.



3. **Resolution inconsistencies:** Most data in the map are at least 30 meter resolution, but some lower resolution data were used when higher resolution data were unavailable. These include the CORINE data at 100 meter resolution, and some countries/regions in the SDPT that included source data at low resolutions (see Appendix D). This resolution inconsistency led to some data artifacts - meaning sharp boundary lines between natural and non-natural areas that are only due to the resolution (Figure 6).

### Natural Lands

- Forests
- Wetland Forests
- Peat Forests
- Short Vegetation
- Wetland Short Vegetation
- Peat Short Vegetation
- Water
- Mangroves
- Bare Ground
- Snow
- Non-Natural




Figure 6. Example of data artifacts present in the map in China due to incorporating the SDPT, which includes source data at 1km resolution for China.

4. **Insufficient data:** Earth observation data is limited in its ability to directly derive information on species composition, structure, and ecological function - the primary elements which define natural ecosystems in the AFI guidance. Some non-natural areas are somewhat easier to delineate with earth observation data though, such as tree plantations, built-up areas and cropland. Natural lands are thus deduced by removing these areas instead of including direct measurements of the definition. Even so, availability of data is not consistent across regions or types of plantations; for example, while there are multiple high-resolution maps of palm oil in some Southeast Asian countries, there are less data available for other palm oil producing regions or other crop types, particularly those grown under shade in agroforestry systems, such as coffee or cocoa. Additionally, certain ecosystems are easier to map than others. The distinction between natural and non-natural short vegetation and bare ground is extremely difficult to identify using earth observation, and wetlands are more difficult to detect during dry periods. As a result of the limitations described above, users should be aware of the following:
  - Natural forests are overestimated in temperate and boreal regions, particularly Europe and Canada. In these regions, it can be challenging to differentiate plantation forests from natural forests, as rotation cycles are often long (greater than 20 years) and therefore not frequently discernable



with earth observation data. Moreover, spatially-explicit data on forest management in these regions is limited. The SDPT v2 primarily uses the Global Forest Management (GFM) data for Europe, Canada, and Russia, as it is currently the best available data for these regions. However, Lesiv et al. (2022) note that planted forests are underestimated in this dataset and as a result, natural forests in the natural lands map are overestimated in these regions. Additionally, the GFM data is representative of the year 2015, and therefore may not include more recently established planted or plantation forests.

- In some regions, small-scale agriculture– particularly mixed/rotational pasture and cropland– is classified as natural forest or short vegetation due to a lack of data that can be used to delineate these areas as non-natural. These heterogeneous agricultural mosaics are not always well-captured in existing cropland data, and due to the rotational nature of these systems, may not be included in existing data on plantations, tree crops, or planted trees.
5. **Tradeoff between global and regional data sources:** The use of regional data sources can be beneficial, especially in overcoming limitations of global data sets and ensuring that local knowledge and conditions are well represented in the map. For example, regional data allowed for better delineation of natural and non-natural short vegetation in South America. It also introduced local definitions, which better account for the unique characteristics of a particular landscape. However, the use of regional data introduces methodological and definitional inconsistencies with areas outside those regions. The inclusion of regional data also introduces artifacts at the edge of the regional data source, often with the regional and global data differing in vegetation type (e.g. bare versus short vegetation, or short vegetation vs forest) but remaining the same natural/non-natural class. Any organization using the natural lands map will need to use caution when comparing performance in supply chains across geographies with different data sources.
  6. **Monitoring challenges:** The natural lands map is only available for the year 2020 and there are currently only limited monitoring systems in place for natural lands. Future mapping efforts should focus on producing dynamic maps which show change across multiple time periods or monitoring systems to evaluate change since the 2020 natural lands baseline. While we are aware of a number of regional land monitoring datasets (PRODES and DETER), a new larger scale alert system (OPERA Disturbance Alerts), and a new grassland monitoring system (Global Pasture Watch) in development, even monitoring in forests, which have had monitoring systems in place for years, has proven to be challenging. Natural forest and short vegetation basemaps will help with a key barrier to uptake of monitoring data by corporate and other actors, which is the delineation of natural and non-natural forests. Another challenge which remains, however, is the lag time associated with deforestation and conversion. While we can now detect the initial forest disturbance quickly, it can take years to understand if the forest then regrew (not deforestation) or if it resulted in a change of land use (deforestation) using only remote sensing data. The land must be cleared and converted to another land use, then detected in a land cover/land use dataset to register deforestation, and this process takes time, often longer than annual reporting cycles.

7. **Equity impact:** The natural lands map and definitions overestimate natural areas, which is intended to protect land from potential conversion. However, a potential drawback of this approach is that areas with less available data to delineate non-natural lands may have a relatively larger overestimation of natural lands. Overestimation may also occur in areas where agricultural production systems do not clearly fall into the natural or non-natural land definitions, such as lower intensity, mixed, and shifting agriculture. Thus this limitation has important social and equity considerations. We recommend that all companies using the natural lands map to set no conversion targets validate the map with high resolution imagery or ground truthing, and engage with the local communities to understand the landscape. We recommend companies also set the SBTN land targets on land footprint reduction and landscape engagement to increase the effectiveness of the no conversion target and minimize unintended negative consequences for the communities where they are producing or sourcing.

## 5. Future Work

The natural lands map version 1 uses the best publicly available data. As new and improved (higher accuracy, finer resolutions, better thematic detail) datasets are published, they could be incorporated into the map as future versions. New data for natural grasslands and planted forests in temperate and boreal regions would be especially valuable to future maps. New global grassland condition and livestock data at 30 meter resolution are expected from [Land & Carbon Lab's Global Grassland Monitoring Consortium](#) in the near future.

Further work in this field is needed to refine the definitions of natural non-forest ecosystems and conversion, and identify and create data capable of monitoring the conversion of natural lands.

## 6. Data Availability

The SBTN Natural Lands Map can be visualized at <https://wri-datalab.earthengine.app/view/sbtn-natural-lands>.

Data are available on Google Earth Engine at [WRI/SBTN/naturalLands/v1\\_1/2020](https://earthengine.google.com/viewer/?layers=WRI%2F%2FSBTN%2FnaturalLands%2Fv1_1%2F2020)

All code is available on Github at the following link: <https://github.com/wri/natural-lands-map/>

The SBTN Natural Lands Map has a Creative Commons Attribution ShareAlike 4.0 [Intergovernmental Organization License](#).

### Acknowledgements

We thank the team at IIASA for validating the natural lands map by collecting over four thousand visually inspected points. The team included Steffen Fritz, Myroslava Lesiv, Katya Perez Guzman, Linda See, Maria Shepaschenko, Ivelina Georgieva, Florian Hofhansl, and Juan Carlos Laso Bayas. Thank you to the WRI staff Lindsey Sloat, Radost Stanimirova, and Ana Reboredo Segovia for assisting in the evaluation of Global Pasture Watch data. We also want to thank our many reviewers for their time and detailed comments that improved the map and technical note, including Sarah Carter, David Gibbs, Viviana Zalles, Kemen Austin, Radost Stanimirova, Megan Critchley, Neil Burgess, Susana Baena, Marcos Rosa, Sveltana Turubanova, Gregory Taff, and the many people who reviewed the map through the SBTN internal review and public consultation. While the contributions from reviewers are greatly appreciated, the approach presented in this report are those of the authors alone.

# Appendix A: MapBiomass Reclassification

We reclassified each MapBiomass dataset to the value that aligned most closely with our natural land classification. For classes that contained both natural forest and natural non-forest vegetation according to the AFI/FAO definitions (such as savannas and woodlands), we overlaid the UMD tree height data (greater than or equal to 5 meters) to delineate natural forest versus natural short vegetation. We then applied a minimum mapping unit of 0.5 hectares and classified any patches that did not meet this threshold as natural short vegetation. In cases where the MapBiomass category was not specific enough to allow differentiation between natural/not natural categories or a specific class on our map, we overlaid the class with UMD Land Cover data to assign categories.

MapBiomass uses a hierarchical classification system, which includes level 1 and 2 classes. Level 1 classes are listed in darker colors in the table below, whereas level 2 classes are listed in lighter colors. Level 2 classes were used to remap to natural lands map classes where available; therefore some classes only include level 2 values in Table A.

The MapBiomass classes present in each dataset are marked with an 'X' in Table A.

Table A: Reclassification table for MapBiomass datasets

MapBiomass Classification															Natural Lands Reclassification		
		Class Number	BRA	Amazon	PER	BOL	COL	VEN	URY	ECU	PRY	CHL	ARG	IDN	Category	Class	Class number
Natural Forest Formation	Natural Forest	1															
	Forest Formation	3	X	X	X	X	X	X	X	X	X	X	X	X	natural	forest	2
	Savanna Formation	4	X	X	X	X		X		X	X		X		natural	forest/ short vegetation*	2/3
	Mangrove	5	X	X	X		X	X		X				X	natural	mangrove	5
	Sandy coastal plain vegetation	49	X					X							natural	forest	2

	Flooded/ wetland forest	6	X	X	X	X	X	X		X	X		X		natural	wetland forest	8
	Sparse woodlands	45											X		natural	forest/ short vegetation*	2/3
Natural Non- Forest Form- ation	<b>Natural Non- Forest Formation</b>	10															
	Wetland	11	X	X	X	X	X	X	X	X	X	X	X		natural	wetland short vegetation	10
	Grassland	12	X	X	X	X	X	X	X	X	X	X	X		natural	short vegetation	3
	Grassland with open vegetation	42													natural	short vegetation	3
	Grassland with closed vegetation	43													natural	short vegetation	3
	Grassland with dispersed vegetation	44													natural	short vegetation	3
	Shrubland	66				X						X			natural	short vegetation	3
	Steppe	63											X		natural	short vegetation	3
	Hypersaline Tidal Flat	32	X		X		X	X							natural	wetland short vegetation	10
	Rocky Outcrop	29	X	X		X	X	X		X		X			natural	bare	6
	Herbaceous Sandbank Vegetation	50	X				X	X							natural	short vegetation	3

	Other non Forest Formations	13	X	X	X	X	X	X		X				X**	natural	short vegetation	3
Farming	Farming	14															
	Pasture	15	X	X	X	X		X	X		X		X		non-natural	short vegetation	15
	Agriculture	18		X	X	X		X					X		non-natural	cropland	12
	Temporary Crop	19							X		X				non-natural	cropland	12
	One crop	57													non-natural	cropland	12
	Multiple crops	58													non-natural	cropland	12
	Soybean	39	X												non-natural	cropland	12
	Sugar cane	20	X												non-natural	cropland	12
	Rice	40	X											X	non-natural	cropland	12
	Cotton	62	X												non-natural	cropland	12
	Other Temporary Crops	41	X												non-natural	cropland	12
	Perennial Crop	36											X		non-natural	cropland	12
	Coffee	46	X												non-natural	tree cover	14
	Citrus	47	X												non-natural	tree cover	14
	Tea	65													non-natural	cropland	14
	Other Perennial Crops	48	X												non-natural	tree cover	14
Oil Palm	35	X	X	X		X							X	non-natural	tree cover	14	

	Forest Plantation	9	X	X	X		X	X	X	X	X	X	X	X	non-natural	tree cover	14
	Mosaic of Uses	21	X	X	X	X	X	X		X		X	X	X	non-natural	cropland	12
Non vegetated area	Non vegetated area	22							X		X		X		natural/non-natural	natural bare/ Built/ Crop***	6/13/12
	Beach, Dune and Sand Spot	23	X	X			X	X		X		X			natural	bare	6
	Urban Area	24	X	X	X	X	X	X		X		X			non-natural	built	13
	Mining	30	X	X	X	X	X	X		X				X	non-natural	built	13
	Salt Flat	61				X						X			natural	bare	6
	Other Non Vegetated Natural Areas	68		X		X	X	X		X					natural	bare	6
	Other Non Vegetated Anthropic Areas	25	X	X	X***	X	X	X		X		X***		X***	natural/ non-natural	natural bare/ Built/ Crop***	6/13/12
Water	Water	26													natural	water	4
	River, Lake and Ocean	33	X	X	X	X	X	X	X	X	X	X	X	X	natural	water	4
	Glacier	34		X	X	X	X	X		X		X	X		natural	snow/ice	7
	Aquaculture	31	X		X		X	X		X				X	non-natural	water	16
	6. Not Observed	27		X	X	X	X		X	X	X	X	X	X	Mask and take global map value		

\*Because the MapBiomas savannas and sparse woodlands classes contain both forest and non-forest natural vegetation according to the AFI/FAO thresholds, we overlaid the UMD tree height data (greater than or equal to 5 meters) to delineate natural forest versus natural short vegetation. We then applied a



minimum mapping unit of 0.5 hectares to the area reclassified as forest, and classified any patches that did not meet this threshold as natural short vegetation.

\*\* MapBiomas Indonesia was the only dataset without a wetlands class, though wetlands are present in Indonesia. First, we reclassified the 'other non-forest formations' class (13) in Indonesia to natural short vegetation. We then overlaid this class with the UMD Land Cover data and reclassified it to natural wetland short vegetation if it overlapped with the UMD wetland short vegetation class.

\*\*\*MapBiomas classes for non-vegetated areas can include non-natural built-up areas and transitional cropland, as well as natural bare land, sand, rock, and other non-vegetated cover. First, we reclassified the 'non-vegetated area' (22) or 'other non-vegetated areas' (25) to natural bare land. We then overlaid this class with the UMD Land Cover data and reclassified it to non-natural built if it overlapped with the UMD built-up class, and overlaid this class with the UMD crop data and reclassified it to crop if it overlapped with the UMD crop class.

\*\*\*\*MapBiomas classes for non-vegetated areas can include non-natural built-up areas and transitional cropland. In order to assign the appropriate land cover class, we reclassified the 'other non-vegetated areas' (25) to non-natural bare land. We then overlaid this class with the UMD Land Cover data and reclassified it to non-natural built if it overlapped with the UMD built-up class, and overlaid this class with the UMD crop data and reclassified it to crop if it overlapped with the UMD crop class.

# Appendix B: South Africa National Land Cover Reclassification

We reclassified the South Africa National Land Cover 2020 data to the value that aligned most closely with our natural land classification. For classes that contained both natural forest and natural non-forest vegetation according to the AFi/FAO definitions (such as mixed classes and woodlands), we overlaid the UMD tree height data (greater than or equal to 5 meters) to delineate natural forest versus natural short vegetation. We then applied a minimum mapping unit of 0.5 hectares and classified any patches that did not meet this threshold as natural short vegetation.

Table B: Reclassification table for the South Africa National Land Cover 2020 Map

South Africa National Land Cover 2020 Classification		Natural Lands Reclassification		
Class number	Name	Category	Class	Class number
1	Contiguous Forest (combined very high, high, medium)	Natural	Forest	2
2	Contiguous Low Forest & Thicket	Natural	Forest/Short vegetation*	2/3
3	Dense Forest & Woodland	Natural	Forest/Short vegetation*	2/3
4	Open Woodland	Natural	Forest/Short vegetation*	2/3
5	Contiguous & Dense Planted Forest	Non-Natural	Forest	14
6	Open & Sparse Planted Forest	Non-Natural	Forest	14
7	Temporary Unplanted Forest	Non-Natural	Forest	14

8	Low Shrubland (other regions)	Natural	Short vegetation	3
9	Low Shrubland (Fynbos)	Natural	Short vegetation	3
10	Low Shrubland (Succulent Karoo)	Natural	Short vegetation	3
11	Low Shrubland (Nama Karoo)	Natural	Short vegetation	3
12	Sparsely Wooded Grassland	Natural	Short vegetation	3
13	Natural Grassland	Natural	Short vegetation	3
14	Natural Rivers	Natural	Water	4
15	Natural Estuaries & Lagoons	Natural	Water	4
16	Natural Ocean	Natural	Water	4
17	Natural Lakes	Natural	Water	4
18	Natural Pans (flooded)	Natural	Water	4
19	Artificial Dams	Non-Natural	Water	16
20	Artificial Sewage Ponds	Non-Natural	Water	16
21	Artificial Flooded Mine Pits	Non-Natural	Water	16
22	Herbaceous Wetlands (currently mapped)	Natural	Wetland short vegetation	10

23	Herbaceous Wetlands (previous mapped extent)	Natural	Wetland short vegetation	10
24	Mangrove Wetlands	Natural	Mangroves	5
25	Natural Rock Surfaces	Natural	Bare	6
26	Dry Pans	Natural	Bare	6
27	Eroded Lands	Natural	Bare	6
28	Sand Dunes (terrestrial)	Natural	Bare	6
29	Coastal Dunes & Beach Sand	Natural	Bare	6
30	Bare Riverbed Material	Natural	Bare	6
31	Other Bare	Natural	Bare	6
32	Cultivated Commercial Permanent Orchards	Non-Natural	Forest	14
33	Cultivated Commercial Permanent Vines	Non-Natural	Cropland	12
34	Cultivated Commercial Sugarcane Pivot Irrigated	Non-Natural	Cropland	12
35	Commercial Permanent (Pineapples)	Non-Natural	Cropland	12
36	Cultivated Commercial Sugarcane Non-Pivot (all other)	Non-Natural	Cropland	12

37	Cultivated Emerging Farmer Sugarcane Non-Pivot (all other)	Non-Natural	Cropland	12
38	Cultivated Commercial Annuals Pivot Irrigated	Non-Natural	Cropland	12
39	Cultivated Commercial Annuals Non-Pivot Irrigated	Non-Natural	Cropland	12
40	Cultivated Commercial Annuals Non-Pivot / Non-Irrigated	Non-Natural	Cropland	12
41	Subsistence Annual Crops	Non-Natural	Cropland	12
42	Fallow Land & Old Fields (Trees)**	Natural	Forest	2
43	Fallow Land & Old Fields (Bush)**	Natural	Short vegetation	3
44	Fallow Land & Old Fields (Grass)**	Natural	Short vegetation	3
45	Fallow Land & Old Fields (Bare)**	Natural	Bare	6
46	Fallow Land & Old Fields (Low Shrub)**	Natural	Short vegetation	3
47	Residential Formal (Tree)	Non-Natural	Forest	14
48	Residential Formal (Bush)	Non-Natural	Short vegetation	15
49	Residential Formal (low veg / grass)	Non-Natural	Short vegetation	15
50	Residential Formal (Bare)	Non-Natural	Built-up	13

51	Residential Informal (Tree)	Non-Natural	Forest	14
52	Residential Informal (Bush)	Non-Natural	Short vegetation	15
53	Residential Informal (low veg / grass)	Non-Natural	Short vegetation	15
54	Residential Informal (Bare)	Non-Natural	Built-up	13
55	Village Scattered	Non-Natural	Built-up	13
56	Village Dense	Non-Natural	Built-up	13
57	Smallholdings (Tree)	Non-Natural	Forest	14
58	Smallholdings (Bush)	Non-Natural	Short vegetation	15
59	Smallholdings (low veg / grass)	Non-Natural	Short vegetation	15
60	Smallholdings (Bare)	Non-Natural	Built-up	13
61	Urban Recreational Fields (Tree)	Non-Natural	Forests	14
62	Urban Recreational Fields (Bush)	Non-Natural	Short vegetation	15
63	Urban Recreational Fields (Grass)	Non-Natural	Short vegetation	15
64	Urban Recreational Fields (Bare)	Non-Natural	Built-up	13
65	Commercial	Non-Natural	Built-up	13
66	Industrial	Non-Natural	Built-up	13

67	Roads & Rail (Major Linear)	Non-Natural	Built-up	13
68	Mines: Surface Infrastructure	Non-Natural	Built-up	13
69	Mines: Extraction Sites: Open Cast & Quarries combined	Non-Natural	Built-up	13
70	Mines: Extraction Sites: Salt Mines	Non-Natural	Built-up	13
71	Mines: Waste (Tailings) & Resource Dumps	Non-Natural	Built-up	13
72	Land-fills	Non-Natural	Built-up	13
73	Fallow Land & Old Fields (wetlands)**	Natural	Wetland short vegetation	10

\*For classes that contained both natural forest and natural non-forest vegetation according to the AFi/FAO definitions, we overlaid the UMD tree height data (greater than or equal to 5 meters) to delineate natural forest versus natural short vegetation. We then applied a minimum mapping unit of 0.5 hectares and classified any patches that did not meet this threshold as natural short vegetation.

\*\*We classified fallow land and old fields as natural, rather than cropland, because the class description states that these are long-term, non-active, previously cultivated lands where the cultivated land unit is no longer detectable, and thus may meet the AFi definition of a regenerated natural ecosystem (which is included in the AFi natural ecosystem definition). These classes were mapped using historical field boundaries from the 1950s-70s.



# Appendix C: LUCAS New Zealand Land Use Reclassification

Table C: Reclassification for the LUCAS New Zealand Land Use dataset

LUCAS NZ Land Use Classification				Natural Lands Reclassification		
ID	Class	Subclass	Subclass ID	Category	Class	Class number
71	Natural Forest	Shrubland	120	Natural	Forest	2
		Tall forest	121	Natural	Forest	2
		Wilding trees	122	Natural	Forest	2
72	Pre-1990 Planted Forest	Unknown	0	Non-Natural	Forest	14
		Pinus radiata	201	Non-Natural	Forest	14
		Douglas fir	202	Non-Natural	Forest	14
		Unspecified exotic species	203	Non-Natural	Forest	14
73	Post-1989 Forest	Wilding Trees	122	Natural	Forest	2
		Pinus radiata	201	Non-Natural	Forest	14
		Douglas Fir	202	Non-Natural	Forest	14
		Unspecified exotic species	203	Non-Natural	Forest	14
		Regenerating natural species	204	Natural	Forest	2

74	Grassland - with woody biomass	Unknown	0	Natural	Short Vegetation	3
75	Grassland - high producing	Unknown	0	Non-Natural	Short Vegetation	15
		Grazed - dairy	502	Non-Natural	Short Vegetation	15
		Grazed - non-dairy	503	Non-Natural	Short Vegetation	15
		Ungrazed	504	Non-Natural	Short Vegetation	15
76	Grassland - low producing	Unknown	0	Natural	Short Vegetation	3
		Grazed - dairy	502	Natural	Short Vegetation	15
		Grazed - non-dairy	503	Natural	Short Vegetation	15
		Ungrazed	504	Natural	Short Vegetation	3
77	Cropland - perennial	Unknown	0	Non-Natural	Cropland	12
78	Cropland - annual	Unknown	0	Non-Natural	Cropland	12
79	Open water	Unknown	0	Natural	Water	4
		Naturally occurring	901	Natural	Water	4
		Human induced	902	Non-Natural	Water	16
80	Vegetated wetland	Unknown	0	Natural	Wetland Short Vegetation	10
		Peat mine	1001	Non-Natural	Wetland Short Vegetation	17
81	Settlements	Unknown	0	Non-Natural	Built-up	13
82	Other	Unknown	0	Natural	Bare / snow/ice *	6/7

\*The LUCAS 'Other' class can include natural bare rock and sand, as well as permanent ice/snow and glaciers. First, we reclassified the Other class as bare and then we overlaid this class with the UMD Land Cover and reclassified it to snow/ice if it overlapped with the UMD snow/ice class.

# Appendix D: SDPT v2.0 Data Sources

For full technical documentation on the methods used to produce the Spatial Database of Planted Trees version 2.0, please see the associated publication. Table D lists the datasets used for each country, as well as new regional datasets incorporated.

Table D: Data sources in SDPT v2.0

Country	Year	Source	Native resolution	Regional - Oil Palm (Descals et al. 2021) 10m	Regional - Rubber (Xiao et al. 2021) 30m	Regional - Orchard (Open Street Map)
Algeria	2015	Lesiv et al. 2022	100 m			x
Angola	-	-		x		x
Argentina	2013	Argentina Ministry of Agroindustry	vector			x
Armenia	2015	Lesiv et al. 2022	100 m			x
Australia	2014-2015	Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES)	vector			x
Azerbaijan	2015	Lesiv et al. 2022	100 m			x
Bangladesh	-	-				x
Belize	2018	Belize Ministry of Agriculture, Fisheries, Forestry, the Environment, Sustainable Development, and Immigration	vector	x		x
Benin	2013	USGS LULC West Africa	2 km	x		x
Bhutan	2015	Lesiv et al. 2022	100 m			x
Bolivia	2015	Lesiv et al. 2022	100 m			x

Botswana	2015	Lesiv et al. 2022	100 m			x
Brazil	2013-2014	Petersen et al. 2016 (Transparent World)	vector	x		x
Brunei	-	-		x		x
Burkina Faso	2013	USGS LULC West Africa	2 km			x
Burundi	-	-		x		x
Cabo Verde	2013	USGS LULC West Africa	2 km			x
Cambodia	2013-2014 / 2015	Petersen et al. 2016 (Transparent World) / Debonne et al. 2019	vector / 1 km	x	x	x
Cameroon	2020	Cameroon Ministry of Forestry and Wildlife, Cameroon Ministry of Forestry and Wildlife/WRI	vector	x		x
Canada	2015	Lesiv et al. 2022	100 m			x
Central African Republic	-	-		x		x
Chile	2014 / 2016	Instituto Forestal de Chile (INFOR), Sistema Informationa de Territorial (SIT) / Zhao et al. 2016	vector / 30m			x
China	2016-2020	Abbasi et al. 2023	1 km			x
Colombia	2013-2014 / 2002-2020	Petersen et al. 2016 (Transparent World) / Instituto Amazónico de Investigaciones Científicas - SINCH	vector / vector	x		x
Congo	-	-		x		x
Costa Rica	2012	Sistema Nacional de Areas de Conservación (SINAC), Fondo Nacional de Financiamiento Forestal (FONAFIFO), Ministerio de Ambiente y Energia (MAE)	5 m	x		x

Cote D'Ivoire	2013-2015/ 2013	WRI/ USGS LULC West Africa	vector/ 2km	x		x
Cuba	2015	Lesiv et al. 2022	100 m			x
Cyprus	2015	Lesiv et al. 2022	100 m			x
Dominican Republic	-	-		x		x
DRC	2013/ 2018	Ministère de L'Environnement, Conservation de la Nature, et Développement Durable (MECNDD), Nature Conservancy and Sustainable Development (MECNDD)/ DRC Ministry of the Environment and Sustainable Development's Forest Atlas	vector/ vector	x		x
Ecuador	2018/ 2020	Ministry of Environment Land Use Map/ Ministry of Agriculture and Livestock	30m / vector	x		x
Egypt	2015	Lesiv et al. 2022	100 m			x
El Salvador	-	-		x		x
Equatorial Guinea	-	-		x		x
Eritrea	2015	Lesiv et al. 2022	100 m			x
Ethiopia	2015	Lesiv et al. 2022	100 m			x
European Union	2015	Lesiv et al. 2022	100 m			x
Fiji	2015	Lesiv et al. 2022	100 m			x
French Guiana	2015	Lesiv et al. 2022	100 m			
Gabon	2013-2015	WRI	vector	x		x

Gambia	2013	USGS LULC West Africa	2 km			x
Ghana	2013-2015/ 2013	WRI/ USGS LULC West Africa	vector/ 2 km	x		x
Guadeloupe	2015	Lesiv et al. 2022	100m			
Guatemala	1998-2020	Guatemala Forestry incentives database, Forestry Development Directorate - INAB 2020	vector	x		x
Guinea	2013	USGS LULC West Africa	2 km	x		x
Guinea-Bissau	2013	USGS LULC West Africa	2 km	x		x
Haiti	2015	Lesiv et al. 2022	100 m			x
Honduras	2013	National Institute of Conservation and Forest Development, Protected Areas, and Wildlife	5 m	x		x
India	2015	Roy et al. 2016	23.5 m	x		x
Indonesia	2013-2014 / 2015/ 2017-2019	Petersen et al. 2016 (Transparent World) / Austin et al. 2017/ Miettinen et al. 2016/ Gaveau et al. 2016/ Condro et al. 2020	vector/ 250 m/ 30 m / 60m/ 30m	x		x
Iran	2015	Lesiv et al. 2022	100 m			x
Iraq	2015	Lesiv et al. 2022	100 m			x
Israel	2015	Lesiv et al. 2022	100 m			x
Jamaica	2015	Lesiv et al. 2022	100 m			x
Japan	2016-2020	Abbasi et al. 2023	1 km			x
Jordan	2015	Lesiv et al. 2022	100 m			x



Kazakhstan	2015	Lesiv et al. 2022	100 m			x
Kenya	2010	Kenya Forest Service	vector			x
Kyrgyzstan	2015	Lesiv et al. 2022	100 m			x
Laos	-	-			x	x
Lebanon	2015	Lesiv et al. 2022	100 m			x
Lesotho	2015	Lesiv et al. 2022	100 m			x
Liberia	2013-2014/ 2013	Petersen et al. 2016 (Transparent World)/ USGS LULC West Africa	vector/ 2 km	x		x
Libya	2015	Lesiv et al. 2022	100 m			x
Madagascar	2015	Lesiv et al. 2022	100 m			x
Malawi	2012/ 2015	Malawi Department of Forestry/ Food and Agriculture Organization (FAO)	vector/ vector			x
Malaysia	2013-2014/ 2015/ 2010/ 2016	Petersen et al. 2016 (Transparent World)/ Gaveau et al. 2016/ Miettinen et al. 2016 / Gunarso et al. 2013/ Xu et al. 2020	vector/ 30 m/ 30 m/ vector/ 100 m	x		x
Mali	2013	USGS LULC West Africa	2 km			x
Mauritania	2013	USGS LULC West Africa	2 km			x
Mexico	2010-2021/ 2018	Dirección General de Gestión Forestal y de Suelos (DGGFS) of Mexico's Ministry of Environment and Natural Resources (Secretaría de Medio Ambiente y Recursos Naturales; SEMARNAT)/ Mexico Conafor Comision Nacional Forestal, INEGI	vector/ vector	x		x

Mongolia	2015	Lesiv et al. 2022	100 m			x
Morocco	2015	Lesiv et al. 2022	100 m			x
Mozambique	2015	Lesiv et al. 2022	100 m			x
Myanmar	2014	Bhagwat et al. (2015)	30 m	x	x	x
Nepal	2015	Ministry of Forest and Soil Conservation	vector			x
New Caledonia	2015	Lesiv et al. 2022	100 m			x
New Zealand	2016	New Zealand Ministry for the Environment LUCAS Land Use	vector			x
Nicaragua	2014	Furumo and Aide (2017)	250 m	x		x
Nigeria	2013-2015/ 2013	WRI/ USGS LULC West Africa	vector/ 2 km	x		x
North Korea	2016-2020	Abbasi et al. 2023	1 km			x
Oman	2015	Lesiv et al. 2022	100 m			x
Pakistan	2015	Pakistan Forestry, Environment and Wildlife Department	vector			x
Palestine	2015	Lesiv et al. 2022	100 m			x
Panama	2021	Panama Ministerio de Ambiente	10 m	x		x
Papua New Guinea	2015/ 2015	Papua New Guinea Forest Authority (PNGFA)/ New Britain Palm Oil Ltd (NBPOL)	2 km/ vector	x		x
Paraguay	2015	Lesiv et al. 2022	100 m			x
Peru	2013-2014	Petersen et al. 2016 (Transparent World)	vector	x		x

Philippines	2003, 2017	National Mapping and Resource Information Authority (NAMRIA)	vector	x		x
Rwanda	2008	Government of Rwanda	vector	x		x
Sao Tome and Principe	-	-		x		x
Senegal	2013	USGS LULC West Africa	2 km			x
Sierra Leone	2013	USGS LULC West Africa	2 km	x		x
Solomon Islands	-	-		x		x
Somalia	2015	Lesiv et al. 2022	100 m			x
South Africa	2020	South Africa Department of Forestry, Fisheries, and the Environment Land Cover Map	20 m			x
South Korea	Unk./ Unk.	Korean Forest Service/ South Korea National Map of Planted Forests (Kim et al. 2009)	vector/ vector			x
South Sudan	2015	Lesiv et al. 2022	100 m			x
Sri Lanka	2013–2015	WRI	vector	x		x
Suriname	2015	Lesiv et al. 2022	100 m			x
Swaziland	2015	Lesiv et al. 2022	100 m			x
Syria	2015	Lesiv et al. 2022	100 m			x
Tajikistan	2015	Lesiv et al. 2022	100 m			x
Tanzania	-	-		x		x
Thailand	2000	Thai Royal Forestry Department	vector	x	x	x
Togo	2013	USGS LULC West Africa	2 km	x		x

Trinidad and Tobago	2007	Helmer et al. 2012	vector			
Tunisia	2015	Lesiv et al. 2022	100 m			x
Uganda	-	-		x		x
United States	2017 / 2014	United States National Agricultural Statistics Service; NASS / WRI analysis based on data from USDA Forest Service (ownership, forest type, timberland extent), US Geological Survey (protected areas), Pan et al. 2011 (stand age)	30 m/ 250m			x
Uruguay	2015/ 2021	Dirección Nacional de Ordenamiento Territorial (DINOT), within Ministerio de Vivienda, Ordenamiento Territorial y Medio Ambiente (MVOTMA) / Uruguay Ministry of Livestock, Agriculture, and Fisheries	vector/ vector			x
Uzbekistan	2015	Lesiv et al. 2022	100 m			x
Vanuatu	-	-		x		x
Venezuela	2014	Furumo and Aide (2017)	250 m	x		x
Vietnam	2016	Government of Vietnam	vector	x	x	x
Zambia	2015	Lesiv et al. 2022	100 m			x
Zimbabwe	2015	Lesiv et al. 2022	100 m			x

\*Year represents the year of source plantation data and not the publication year.

# Appendix E: Data Source Licenses

Dataset name	Category	Original data license
UMD GLAD Land Cover	global	Creative Commons Attribution 4.0 International License
ESA WorldCover	global	Creative Commons Attribution 4.0 International License
USGS Cropland	global	Public domain
ESA WorldCereal	global	Creative Commons Attribution 4.0 International License
Spatial Database of Planted Trees	global	Creative Commons Attribution 4.0 International License
Global Pasture Watch	global	MIT License
IIASA Global Mining Polygons	global	Creative Commons Attribution - ShareAlike 4.0 International
Global Mining Footprint (Tang et al. 2023)	global	Creative Commons Attribution 4.0 International License
Global Alluvial Mining (Dethier et al. 2023)	global	Creative Commons Attribution 4.0 International License
Intact Forest Landscapes	global	Creative Commons Attribution 4.0 International License
Global Mangrove Watch	global	Creative Commons Attribution 4.0 International License
Global closed-canopy coconut	global	Creative Commons Attribution 4.0 International License
Mapbiomas - Brazil	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mabiomas - Amazon	regional	"public, open and free by simply referring the source"
Mapbiomas - Indonesia	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Peru	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Bolivia	regional	"public, open and free by simply referring the source"

Mapbiomas - Colombia	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Venezuela	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Uruguay	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Ecuador	regional	Creative Commons Attribution 4.0 International License
Mapbiomas - Paraguay	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Chile	regional	Creative Commons Attribution - ShareAlike 4.0 International
Mapbiomas - Argentina	regional	Creative Commons Attribution - ShareAlike 4.0 International
USGS National Land Cover Database	regional	Public domain
South Africa National Land Cover	regional	Creative Commons Attribution 4.0 International License
ETH/EcoVision Cocoa Map	regional	MIT License
LUCAS NZ Land Use Map	regional	Creative Commons Attribution 4.0 International License
CORINE Land Cover	regional	"full free and open data policy, which allows the use of the product(s) also for any commercial purpose."
European Primary Forest Database	regional	Creative Commons Attribution 4.0 International License
DEA Africa Cropland	regional	Creative Commons Attribution 4.0 International License

# References

- Accountability Framework Initiative (AFi), 2024. *Accountability Framework version 1.0*. <https://accountability-framework.org/>
- Accountability Framework Initiative (AFi). 2023. *Operational Guidance on Applying the Definitions Related to Deforestation, Conversion, and Protection of Ecosystems*.
- Baeza, S., E. Vélez-Martin, D. De Abelleira, S. Banchemo, F. Gallego, J. Schirmbeck, S. Veron, et al. 2022. "Two Decades of Land Cover Mapping in the Río de La Plata Grassland Region: The MapBiomass Pampa Initiative." *Remote Sensing Applications: Society and Environment* 28 (November): 100834. doi:[10.1016/j.rsase.2022.100834](https://doi.org/10.1016/j.rsase.2022.100834).
- Barnosky, A.D., N. Matzke, S. Tomiya, G.O.U. Wogan, B. Swartz, T.B. Quental, C. Marshall, J.L. McGuire, et al. 2011. "Has the Earth's sixth mass extinction already arrived?" *Nature* 471: 51-57.
- Bastin, J.-F., N. Berraahmouni, A. Grainger, D. Maniatis, D. Mollicone, R. Moore, C. Patriarca, et al. 2017. "The Extent of Forest in Dryland Biomes." *Science* 356 (6338): 635-38. doi:[10.1126/science.aam6527](https://doi.org/10.1126/science.aam6527).
- Bunting, P., A. Rosenqvist, L. Hilarides, R.M. Lucas, N. Thomas, T. Tadono, T.A. Worthington, M. Spalding, N.J. Murray, and L.M. Rebelo. 2022. "Global Mangrove Extent Change 1996-2020: Global Mangrove Watch Version 3.0." *Remote Sensing* 14 (3657). <https://doi.org/10.3390/rs14153657>
- Clewley, D., P. Bunting, J. Shepherd, S. Gillingham, N. Flood, J. Dymond, R. Lucas, J. Armston, and M. Moghaddam. 2014. "A Python-Based Open Source System for Geographic Object-Based Image Analysis (GEOBIA) Utilizing Raster Attribute Tables." *Remote Sensing* 6 (7): 6111-35. doi:[10.3390/rs6076111](https://doi.org/10.3390/rs6076111).
- Cover, C.L., 2018. European Union, Copernicus Land Monitoring Service. Eur. Environ. Agency.
- Crezee, B., G.C. Dargie, C.E.N. Ewango, E.T.A. Mitchard, O. Emba, J. Kanyama, P. Bola, J.B.N. Ndjango, N.T. Girkin, Y.E. Bocko, S.A. Ifo, W. Hubau, D. Seidensticker, R. Batumike, G. Imani, A. Cuni-Sanchez, C.A. Kiahtipes, J. Lebamba, H.P. Wotzka, H. Bean, T.R. Baker, A.J. Baird, A. Boom, P.J. Morris, S.E. Page, I.T. Lawson, and S.L. Lewis. "Mapping peat thickness and carbon stocks of the central Congo Basin using field data." *Nature Geoscience* 15: 639-644. <https://www.nature.com/articles/s41561-022-00966-7>
- de Abelleira, D., S. Verón, S. Banchemo, M. Iturralde Elortegui, S. Valiente, O. Puig, F. Murray, J.P. Martini, K. Zelaya, D. Maidana, N. Varlamoff, J. Peiretti, P. Benedetti, J. Portillo, M. Melilli, E. Maidana, Y. Goytía, V. Sapino, A. Regonat, F. Cracogna, C. Espíndola, M. Parodi, F. Colombo, J. Scarel, J. Ayala, L. Martins, M. Basanta, A. Rausch, G. Almada, L. Boero, J. Calcha, A. Chiavassa, M. Lopez de Sabando, D. Kurtz, F. Marini, S. Sarrailhe, M. Petek, T. Propato, A. Ferraina, A. Pezzola, C. Winschel, S. Muñoz, J. Mesa, J. Volante. 2021. "Mapa Nacional de Cultivos Campaña 2020/2021." Instituto Nacional de Tecnología Agropecuaria.
- de Abelleira, D., S. Verón, S. Banchemo, J. Mosciaro, A. Franzoni, M. Boasso, S. Valiente, O. Puig, Y. Goytía, M.R. Iturralde Elortegui, D. Maidana, J.P. Martini, F. Murray, F. Marini, T. Propato, A. Ferraina, C. Gomez, S. Sarrailhe, M. Petek, J.C. Mesa, D. Kurtz, R. Perucca, P. Benedetti, S. Muñoz, J. Volante. 2020. "Mapa Nacional de Cultivos Campaña 2019/2020." Instituto Nacional de Tecnología Agropecuaria.
- Descals, A., S. Wich, E. Meijaard, D. L. A. Gaveau, S. Peedell, and Z. Szantoi. 2021. "High-resolution global map of smallholder and industrial closed-canopy oil palm plantations." *Earth System Science Data* 13(3): 1211-1231. <https://doi.org/10.5194/essd-13-1211-2021>
- Descals, A., S. Wich, Z. Szantoi, M.J. Struebig, R. Dennis, Z. Hatton, T. Ariffin, N. Unus, D.L.A.



- Gaveau, and E. Meijaard. 2023. "High-Resolution Global Map of Closed-Canopy Coconut Palm." *Earth System Science Data* 15 (9): 3991–4010. doi:[10.5194/essd-15-3991-2023](https://doi.org/10.5194/essd-15-3991-2023).
- Dethier, E.N., M. Silman, J.D. Leiva, S. Alqahtani, L.E. Fernandez, P. Pauca, S. Çamalan, et al. 2023. "A Global Rise in Alluvial Mining Increases Sediment Load in Tropical Rivers." *Nature* 620 (7975): 787–93. doi:10.1038/s41586-023-06309-9.
- Digital Earth Africa. 2022. *Cropland extent maps for Africa*. Retrieved 04/2024 from Google Earth Engine. [https://docs.digitalearthafrika.org/en/latest/data\\_specs/Cropland\\_extent\\_specs.html](https://docs.digitalearthafrika.org/en/latest/data_specs/Cropland_extent_specs.html)
- Forest Europe. 2020. "State of Europe's Forest 2020". Ministerial Conference on the Protection of Forests in Europe Liaison Unit Bratislava. [https://foresteurope.org/wp-content/uploads/2016/08/SoEF\\_2020.pdf](https://foresteurope.org/wp-content/uploads/2016/08/SoEF_2020.pdf)
- Gorelick, N., M. Hancher, M. Dixon, S. Llyushchenko, D. Thau, and R. Moore. 2017. "Google earth engine: Planetary-scale geospatial analysis for everyone." *Remote Sensing of Environment*. 202: 18–27. <https://www.sciencedirect.com/science/article/pii/S0034425717302900>
- Gosling, J., M.I. Jones, A. Arnell, J.E.M. Watson, O. Venter, A.C. Baquero, and N.D. Burgess. 2020. "A Global Mapping Template for Natural and Modified Habitat across Terrestrial Earth." *Biological Conservation* 250 (October): 108674. doi:[10.1016/j.biocon.2020.108674](https://doi.org/10.1016/j.biocon.2020.108674).
- Gumbrecht, T., R.M. Roman-Cuesta, L. Verchot, M. Herold, F. Wittman, E. Householder, N. Herold, and D. Murdiyarsa. 2017. "An expert system model for mapping tropical wetlands and peatlands reveals South America as the largest contributor". *Global Change Biology* 23 (9): 3581–3599. <https://onlinelibrary.wiley.com/doi/full/10.1111/gcb.13689>
- Hansen, M.C., P.V. Potapov, A.H. Pickens, A. Tyukavina, A. Hernandez-Serna, V. Zalles, S. Turubanova, I. Kommareddy, S.V. Stehman, and X-P Song. 2022. "Global land use extent and dispersion within natural land cover using Landsat data." *Environmental Research Letters* 17 (3). <https://doi.org/10.1088/1748-9326/ac46ec>
- Harris, N.L., E.D. Goldman, and S. Gibbes. 2019. "Spatial Database of Planted Trees Version 1.0." Technical Note. Washington, DC: World Resources Institute. <https://www.wri.org/publication/spatialdatabase-planted-trees>.
- Harris, N.L., D.A. Gibbs, A. Baccini, R.A. Birdsey, S. de Bruin, M. Farina, L. Fatoyinbo, M.C. Hansen, M. Herold, R.A. Houghton, P.V. Potapov, D. Requena Suarez, R.M. Roman-Cuesta, S.S. Saatchi, C.M. Slay, S.A. Turubanova, A. Tyukavina. 2021. "Global maps of twenty-first century forest carbon fluxes". *Nature Climate Change* 11: 234–240. <https://www.nature.com/articles/s41558-020-00976-6>
- Hastie, A., E.N. Honorio Coronado, J. Reyna, E.T.A. Mitchard, C.M. Akesson, T.R. Baker, L.E.S. Cole, C.J. Cordova Oroche, G. Dargie, N. Davila, E.C. De Grandi, J. Del Aguila, D. Del Castillo Torres, R. De La Cruz Paiva, F.C. Draper, G. Flores, J. Grandez, K. Hergoualc'h, J.E. Householder, J.P. Janovec, O. Lahteenoja, D. Reyna, P. Rodriguez-Veiga, K.H. Roucoux, M. Tobler, C.E. Wheeler, M. Williams, and I.T. Lawson. 2022. "Risks to carbon storage from land-use change revealed by peat thickness maps of Peru". *Nature Geoscience* 15: 369–374. <https://www.nature.com/articles/s41561-022-00923-4>
- Hill, S.L.L., J. Fajardo, C. Maney, M. Harfoot, M. Harrison, D. Guaras, M. Jones, M.J. Oliva, et al. In Review. "The Ecosystem Integrity Index: a novel measure of terrestrial ecosystem integrity with global coverage." *bioRxiv*. <https://doi.org/10.1101/2022.08.21.504707>
- IPBES. 2019. "Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services." S. Díaz, J. Settele, E.S. Brondízio, H.T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, et al. (Eds.). Bonn, Germany: IPBES secretariat. <https://doi.org/10.5281/zenodo.3553579>

IPCC. 2019. "Climate Change and Land, an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems." P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Portner, D.C. Roberts, P. Zhai, R. Slade, et al. (Eds.). Cambridge, United Kingdom and New York, USA: Cambridge University Press.

[https://www.ipcc.ch/site/assets/uploads/sites/4/2022/11/SRCCL\\_Full\\_Report.pdf](https://www.ipcc.ch/site/assets/uploads/sites/4/2022/11/SRCCL_Full_Report.pdf)

Kalischek, N., N. Lang, C. Renier, R.C. Daudt, T. Addoah, W. Thompson, W.J. Blaser-Hart, R. Garrett, K. Schindler, and J.D. Wegner. 2023. "Cocoa Plantations Are Associated with Deforestation in Côte d'Ivoire and Ghana." *Nature Food* 4 (5): 384–93. doi:[10.1038/s43016-023-00751-8](https://doi.org/10.1038/s43016-023-00751-8).

Lesiv, M., D. Shchepashchenko, M. Buchhorn, L. See, M. Dürauer, I. Georgieva, M. Jung, F. Hofhansl, et al. 2022. "Global forest management data for 2015 at a 100m resolution." *Scientific Data* 9 (199). <https://doi.org/10.1038/s41597-022-01332-3>

Ministry for the Environment. 2020. LUCAS NZ land use map 1990 2008 2012 2016 2020 v003. Ministry for the Environment, Wellington

MapBiomias Amazonia Project - Collection 6.0 of Amazonian Annual Land Cover & Land Use Map Series, accessed 01/2025 through Google Earth Engine.

MapBiomias Argentina Project - Collection 1.0 of the Annual Series of Land Cover and Land Use Maps of Argentina, acquired 7/2024 through Google Earth Engine.

MapBiomias Bolivia Project - Collection 2.0 of the Annual Land Cover and Use Maps of Bolivia, acquired on 1/2025 through Google Earth Engine.

MapBiomias Chile Project - Collection 1.0 of the Annual Coverage and Land Use Series, accessed 05/2024 through Google Earth Engine.

MapBiomias Colombia Project - Collection 2.0 of the Annual Series of Cover and Land Use Maps of Colombia, accessed 1/2025 through Google Earth Engine.

MapBiomias Ecuador Project - Collection 2.0 of the Annual Land Cover and Land Use Maps of Ecuador, acquired on 1/2025 through Google Earth Engine.

MapBiomias Indonesia - Collection 2.0 time-series maps of land-use and land-cover, accessed 04/2024 through Google Earth Engine.

MapBiomias Paraguay - Collection 1.0 of the Annual Series of Land Use and Land Cover Maps, acquired on 4/2024 through Google Earth Engine. MapBiomias Peru Project - Collection 2.0 of the Annual Land Cover and Land Use Series for Peru, accessed on 04/2024 through Google Earth Engine.

MapBiomias Uruguay - Collection 2.0 of the Annual Series of Land Use and Land Cover Maps, acquired on 1/2025 through Google Earth Engine.

MapBiomias Venezuela Project - Collection 2.0 of the Annual Series of Land Use and Land Cover Maps of Venezuela, accessed 1/2025 through Google Earth Engine.

Maus, V., Giljum, S., da Silva, D.M. et al. 2022. An update on global mining land use. *Sci Data* 9, 433 <https://doi.org/10.1038/s41597-022-01547-4>

Miettinen, J., C. Shi, and S.C. Liew. 2016. "Land cover distribution in the peatlands of Peninsular Malaysia, Sumatra and Borneo in 2015 with changes since 1990". *Global Ecology and Conservation* 6: 67–78.

<https://www.sciencedirect.com/science/article/pii/S2351989415300470>

Parente, L., Sloat, L., Mesquita, V. et al. Annual 30-m maps of global grassland class and extent (2000–2022) based on spatiotemporal Machine Learning. *Sci Data* 11, 1303 (2024). <https://doi.org/10.1038/s41597-024-04139-6>

Potapov, P., M.C. Hansen, L. Laestadius, S. Turubanova, A. Yaroshenko, C. Thies, W. Smith, I.

Zhuravleva, A. Komarova, S. Minnemeyer, and E. Esipova. 2017. "The last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to 2013." *Science Advances* 3 (1). <https://www.science.org/doi/10.1126/sciadv.1600821>

Richter, J., Goldman, E.D., Harris, N., Gibbs, D., Rose, M., Peyer, S., Richardson, S., Velappan, H.J. 2024. "Spatial Database of Planted Trees Version 2.0." Technical Note. Washington, DC: World Resources Institute. doi: <https://doi.org/10.46830/writn.23.00073>

Sabatini, F.M., H. Bluhm, Z. Kun, D. Aksenov, J.A. Atauri, E. Buchwald, S. Burrascano, E. Cateau, et al. 2021. "European primary forest database v2.0." *Scientific Data* 8 (220). <https://doi.org/10.1038/s41597-021-00988-7>

South Africa Department of Environment, Forestry, and Fisheries (DEFF). 2021. "South African National Land-Cover 2020 Dataset". Downloaded 02/2023 from [https://egis.environment.gov.za/data\\_egis/data\\_download/current](https://egis.environment.gov.za/data_egis/data_download/current).

Souza, Carlos M., Jr., Julia Z. Shimbo, Marcos R. Rosa, Leandro L. Parente, Ane A. Alencar, Bernardo F. T. Rudorff, Heinrich Hasenack, Marcelo Matsumoto, Laerte G. Ferreira, Pedro W. M. Souza-Filho, Sergio W. de Oliveira, Washington F. Rocha, Antônio V. Fonseca, Camila B. Marques, Cesar G. Diniz, Diego Costa, Dyeden Monteiro, Eduardo R. Rosa, Eduardo Vélez-Martin, Eliseu J. Weber, Felipe E. B. Lenti, Fernando F. Paternost, Frans G. C. Pareyn, João V. Siqueira, José L. Viera, Luiz C. Ferreira Neto, Marciano M. Saraiva, Marcio H. Sales, Moises P. G. Salgado, Rodrigo Vasconcelos, Soltan Galano, Vinicius V. Mesquita, and Tasso Azevedo. 2020. "Reconstructing Three Decades of Land Use and Land Cover Changes in Brazilian Biomes with Landsat Archive and Earth Engine" *Remote Sensing* 12, no. 17: 2735. <https://doi.org/10.3390/rs12172735>.

Tang, L., and T.T. Werner. 2023. "Global Mining Footprint Mapped from High-Resolution Satellite Imagery." *Communications Earth & Environment* 4 (1): 1–12. doi:10.1038/s43247-023-00805-6.

Taylor, R., A. Rosenbarger, L. Samberg, J. Milder. "It's Time for Supply Chains to Go Conversion Free — For the Climate's Sake." World Resources Institute. <https://www.landcarbonlab.org/news-updates/supply-chains-zero-land-conversion-climate>

Thenkabail, P.S., P.G. Teluguntla, J. Xiong, A. Oliphant, R.G. Congalton, M. Ozdogan, M.K. Gumma, et al. 2021. "Global Cropland-Extent Product at 30-m Resolution (GCEP30) Derived from Landsat Satellite Time-Series Data for the Year 2015 Using Multiple Machine-Learning Algorithms on Google Earth Engine Cloud." Report 1868. Professional Paper. Reston, VA. USGS Publications Warehouse. doi:[10.3133/pp1868](https://doi.org/10.3133/pp1868).

Tolan, J., H.-I. Yang, B. Nosarzewski, G. Couairon, H.V. Vo, J. Brandt, J. Spore, et al. 2024. "Very High Resolution Canopy Height Maps from RGB Imagery Using Self-Supervised Vision Transformer and Convolutional Decoder Trained on Aerial Lidar." *Remote Sensing of Environment* 300 (January): 113888. doi:10.1016/j.rse.2023.113888.

U.S. Geological Survey (USGS), 2024, Annual NLCD Collection 1 Science Products: U.S. Geological Survey data release, <https://doi.org/10.5066/P94UXNTS>.

Van Tricht, K., Degerickx, J., Gilliams, S., Zanaga, D., Battude, M., Grosu, A., Brombacher, J., Lesiv, M., Bayas, J. C. L., Karanam, S., Fritz, S., Becker-Reshef, I., Franch, B., Mollà-Bononad, B., Boogaard, H., Pratihast, A. K., Koetz, B., and Szantoi, Z.: WorldCereal: a dynamic open-source system for global-scale, seasonal, and reproducible crop and irrigation mapping, *Earth Syst. Sci. Data*, 15, 5491–5515, <https://doi.org/10.5194/essd-15-5491-2023>, 2023.

Venter, O., E.W. Sanderson, A. Magrath, J.R. Allan, J. Beher, K.R. Jones, H.P. Possingham, et al. 2016. "Sixteen Years of Change in the Global Terrestrial Human Footprint and Implications for Biodiversity Conservation." *Nature Communications* 7 (1): 12558. doi:[10.1038/ncomms12558](https://doi.org/10.1038/ncomms12558).

- WWF. 2022. "Living Planet Report 2022 - Building a nature-positive society." R.E.A. Almond, M. Grooten, D. Juffe Bignoli, and T. Petersen (Eds). Gland, Switzerland: World Wildlife Fund. [https://wwflpr.awsassets.panda.org/downloads/lpr\\_2022\\_full\\_report.pdf](https://wwflpr.awsassets.panda.org/downloads/lpr_2022_full_report.pdf)
- Xiao, C., P. Li, Z. Feng, Y. Yang, Z. You, Y. Lin, and X. Zhang. 2021. "Latest 30-m Map of Mature Rubber Plantations in Mainland Southeast Asia and Yunnan Province of China: Spatial Patterns and Geographical Characteristics." *Progress in Physical Geography: Earth and Environment* 45 (5): 736–56. doi:[10.1177/0309133320983746](https://doi.org/10.1177/0309133320983746).
- Xu, J., P.J. Morris, J. Liu, and J. Holden. 2018. "PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis". *CATENA* 160: 134–140. <https://www.sciencedirect.com/science/article/abs/pii/S0341816217303004#ec0005>
- Xu, P. N.E. Tsendbazar, M. Herold, S. de Bruin, M. Koopmans, T. Birch, S. Carter, S. Fritz, M. Lesiv, E. Mazur, A. Pickens, P. Potapov, F. Stolle, A. Tyukavina, R. Van De Kerchove, D. Zanaga. 2024. "Comparative validation of recent 10m-resolution global land cover maps." *Remote Sensing of Environment*, Volume 311, 114316. <https://doi.org/10.1016/j.rse.2024.114316>.
- Zanaga, D., R. Van De Kerchove, W. De Keersmaecker, N. Souverijns, C. Brockmann, R. Quast, J. Wevers, A. Grosu, et al. 2021. "ESA WorldCover 10 m 2020 v100." <https://doi.org/10.5281/zenodo.5571936>



**SCIENCE BASED TARGETS NETWORK**  
GLOBAL COMMONS ALLIANCE